Asthma plan Emergency Department discharge	Patient Details
Your reliever is	
Your preventer is	
Your steroid is	Your peak flow at discharge L/min
Medication	

Day 1	Date		
Reliever		puffs every	hours
Preventer		puffs	times per day
Steroid		mg or	ml daily

Date		
	puffs every	hours
	puffs	times per day
	mg or	ml daily
	Date	puffs every puffs

Day 3	Date		
Reliever		puffs every	hours
Preventer		puffs	times per day
Steroid		mg or	ml daily
Steroid		1119 01	nn dany

Day 4	Date		
Reliever		puffs every	hours
Preventer		puffs	times per day
Steroid		mg or	ml daily

Day 5	
Date	If symptoms are still present please see your General Practitioner and review your Asthma Action Plan

When do I return to the Emergency Department for further review?

If you are using your reliever more frequently than 2 hourly
If there are no signs of improvement or you are feeling worse after 48 hours





What symptoms to look out for

When well



You will

- have no night time wheezing, coughing or chest tightness
- only occasional wheezing, coughing or chest tightness during the day
- need reliever medication only occasionally or before exercise
- do your usual activities without getting asthma symptoms

This is your baseline

When not well



You will

- have increasing night-time wheezing, coughing or chest tightness
- have symptoms regularly in the morning when you wake up
- have a need for extra doses of reliever medication
- have symptoms which interfere with exercise and your usual activities

This is an asthma flare-up

When unwell



You are having an asthma attack when

- you have worsening wheezing, coughing, chest tightness or shortness of breath
- you are waking often at night with asthma symptoms
- you need to use your reliever medication again within 2 hours

This is a severe asthma attack – SEEK MEDICAL ASSISTANCE

Responding to a life threatening asthma attack

If someone near you is having an asthma attack:

- Be calm and reassuring
- Leave them in their most comfortable position
- Dial 000 for an ambulance
- Give 4 puffs of the reliever every 4 minutes via spacer until the ambulance arrives