

In brief

Cerebral venous sinus thrombosis after AstraZeneca vaccination

30 April 2021

Background

- In recent weeks there have been concerns about [blood clots](#) occurring in patients after they were given the AstraZeneca vaccine.(1) Most reports involved [women under 55 years](#).(2)
- These reports included [18 cases](#) of cerebral venous sinus thrombosis (as of 17 March 2021).(3)
- Cerebral venous sinus thrombosis (CVST) refers to the presence of a blood clot in the dural venous sinuses, which drain blood from the brain. Symptoms may include: headache, abnormal vision, any of the symptoms of stroke, such as weakness of the face and limbs on one side of the body and seizures.(4)
- CVST is rare, occurring at a rate of between two and five people per million.(4)
- Cerebral venous sinus thrombosis (CVST) can be a complication of COVID-19. A case series of 14 patients noted most received [anticoagulation](#) (91.7%) and a mortality rate of 45.5%.(5, 6)
- The main treatment for CVST is anticoagulation. There are, however, concerns that heparin is contraindicated in the rare cases of CVST following vaccination (particularly with AstraZeneca vaccine).(7)
- It has been proposed in [Germany](#) and [Norway](#) that post-AstraZeneca vaccination CVST may be similar to a syndrome known as heparin-induced thrombocytopenia or HIT (sometimes referred to as heparin-induced thrombotic thrombocytopenia or HITT).(8, 9)
- HIT is characterised by an anti-platelet factor 4 [antibody response](#) – leading to platelet consumption and thrombosis.(10)
- Notably, HIT features [high thrombotic risk](#) despite only mild to moderate thrombocytopenia. For example, the median platelet count nadir in HIT is approximately 55 to 70 × 10⁹/L, with a high proportion of patients (~30-50%) with platelet count nadirs >100 × 10⁹/L or even >150 × 10⁹/L developing thrombotic events.(11)
- It is diagnosed by a [HITTS screen](#).(12)
- The [standard anticoagulants](#) used in HIT are argatroban (not currently registered in Australia), and bivalirudin.(10)
- The UK Medicines and Healthcare products Regulatory Agency (MHRA) is undertaking a [detailed review](#) of the five cases of CVST with low blood platelets that occurred in the UK, and also notes that these events can occur naturally.(13)

Note: While early minor headache is common with COVID-19 vaccines, it is delayed onset (within days 4-14) of persisting headache, dizziness or visual changes in particular, that indicate further investigation of CVST; with platelet count and film, then antibodies and imaging.

Current Guidance

- [World Health Organisation](#) (19 March 2021): AstraZeneca COVID-19 vaccine continues to have a positive benefit-risk profile and available data does not suggest any overall increase in clotting conditions. There have been very rare reports of thromboembolic events in combination with thrombocytopenia, such as CVST in Europe, however no causal relationship has been established. Education should be provided to health-care professionals and people being vaccinated to recognise the signs and symptoms of all serious adverse events and ongoing monitoring and investigation of adverse events should be done.(14)
- [Australian Government Department of Health](#) (25 March 2021): Benefits of the COVID-19 vaccine far outweigh this potential risk in people with history of clotting conditions. However, vaccination with any COVID-19 vaccine should be deferred for people who have a history of CVST and/or heparin-induced thrombocytopenia. This is only a precautionary measure until further information from ongoing investigations is available.(15)
- [Australian Therapeutic Goods Administration](#) (19 March 2021): The TGA has not received any reports of blood clots following administration of the AstraZeneca COVID-19 vaccine in Australia. It advises that people continue to get the AstraZeneca vaccine when eligible.(13)
- [Medicines and Healthcare products Regulatory Agency](#) (18 March 2021): UK advice remains that the benefits of the vaccines against COVID-19 continue to outweigh any risks and that the public should continue to get their vaccine when invited to do so.(16)
- [International Society on Thrombosis and Haemostasis \(ISTH\)](#) (12 March 2021): Recommends that all eligible adults continue to receive their COVID-19 vaccinations even for patients with a history of blood clots or for those taking blood thinning medications.(17)
- [HSE National Immunisation Office](#) (19 March 2021): Those receiving long term anticoagulation with either warfarin or heparin are not considered to be at higher risk of bleeding complications following immunisation.(18)
- [NHS](#) (last updated 12 Feb 2021): The vaccine can be given intramuscularly to individuals with a bleeding disorder. If the patient is receiving regular treatment to reduce bleeding (for example, patients with haemophilia) vaccine administration can be scheduled to occur shortly after this treatment is given.(19)

To inform this brief, PubMed and targeted website and Google searches were completed on 25 March 2021. The Critical Intelligence Unit maintains a living [evidence table on COVID-19 vaccines](#).

References

1. COVID-19 Critical Intelligence Unit. Emerging evidence on AstraZeneca vaccine and blood clots. 2021;[Internet]. [cited 2021 April 4]. Available at: https://aci.health.nsw.gov.au/_data/assets/pdf_file/0006/639114/Evidence-Check-AstraZeneca-blood-clots.pdf.
2. Therapeutic Goods Administration. Specific clotting condition reported after COVID-19 vaccination. TGA [internet]. [cited 2021 April 9]. Available at: <https://www.tga.gov.au/media-release/specific-clotting-condition-reported-after-covid-19-vaccination>.
3. Mahase E. Covid-19: AstraZeneca vaccine is not linked to increased risk of blood clots, finds European Medicine Agency. BMJ. 2021;372:n774.
4. Devasagayam S, Wyatt B, Leyden J, et al. Cerebral venous sinus thrombosis incidence is higher than previously thought: a retrospective population-based study. Stroke. 2016;47(9):2180-2.

5. Abouhashem S, Eldawoody H, Taha MM. Cerebral venous sinus thrombosis in patients with COVID-19 infection. *Interdisciplinary neurosurgery : Advanced techniques and case management*. 2021;24:101091.
6. Tu TM, Goh C, Tan YK, et al. Cerebral venous thrombosis in patients with COVID-19 infection: a case series and systematic review. *Journal of stroke and cerebrovascular diseases : the official journal of National Stroke Association*. 2020;29(12):105379.
7. Department of health. Updated ATAGI statement for healthcare providers on a specific clotting condition being reported after COVID-19 vaccination. Australian Government [Internet] [cited 2021 April 14] Available at: <https://www.health.gov.au/news/atagi-statement-healthcare-providers-specific-clotting-condition-reported-after-covid-19-vaccination>. 2021.
8. Anderson M. European scientists say they've found link between AstraZeneca's vaccine and blood clots. *Becker's hospital review* [internet] [cited 2021 april 14] Available at: <https://www.beckershospitalreview.com/pharmacy/european-scientists-say-they-ve-found-link-between-astrazeneca-s-vaccine-and-blood-clots.html>. 2021.
9. Steenhuisen J. Scientists probe new theories on whether AstraZeneca shot linked to blood clots. *Reuters* [internet] [cited 2021 April 14] Available at: <https://www.reuters.com/article/us-health-coronavirus-vaccine-idUSKBN2BC01M>. 2021.
10. Gonzales M, Pipalia A, Weil A. Refractory heparin-induced thrombocytopenia with cerebral venous sinus thrombosis treated with IVIg, steroids, and a combination of anticoagulants: a case report. *Journal of investigative medicine high impact case reports*. 2019;7:2324709619832324.
11. Warkentin TE, Kaatz S. COVID-19 versus HIT hypercoagulability. *Thrombosis research*. 2020;196:38-51.
12. RCPA. Heparin-induced thrombocytopenia investigation. The Royal College of Pathologists of Australia [internet] [cited 2021 April 14] Available at: <https://www.rcpa.edu.au/Manuals/RCPA-Manual/Pathology-Tests/H/Heparin-induced-thrombocytopenia-investigation>.
13. Department of Health. AstraZeneca ChAdOx1-S COVID-19 vaccine. 2021; Australian Government [internet]. [cited 2021 April 14]. Available at: <https://www.tga.gov.au/alert/astrazeneca-chadox1-s-covid-19-vaccine>.
14. World Health Organisation. Statement of the WHO Global Advisory Committee on Vaccine Safety (GACVS) COVID-19 subcommittee on safety signals related to the AstraZeneca COVID-19 vaccine. 2021; WHO [internet]. [cited 2021 April 14]. Available at: [https://www.who.int/news/item/19-03-2021-statement-of-the-who-global-advisory-committee-on-vaccine-safety-\(gacvs\)-covid-19-subcommittee-on-safety-signals-related-to-the-astrazeneca-covid-19-vaccine](https://www.who.int/news/item/19-03-2021-statement-of-the-who-global-advisory-committee-on-vaccine-safety-(gacvs)-covid-19-subcommittee-on-safety-signals-related-to-the-astrazeneca-covid-19-vaccine).
15. Department of Health. ATAGI statement for health care providers on suitability of COVID-19 vaccination in people with history of clotting conditions. 2021; Australian Government [internet]. [cited 2021 April 14]. Available at: <https://www.health.gov.au/news/atagi-statement-for-health-care-providers-on-suitability-of-covid-19-vaccination-in-people-with-history-of-clotting-conditions>.
16. UK Government. Government response. UK regulator confirms that people should continue to receive the COVID-19 vaccine AstraZeneca. 2021; [internet]. [cited 2021 April 14]. Available at: <https://www.gov.uk/government/news/uk-regulator-confirms-that-people-should-continue-to-receive-the-covid-19-vaccine-astrazeneca>.
17. Masten A. ISTH statement of Astrazeneca COVID-19 vaccine and thrombosis. 2021; ISTH [internet]. [cited 2021 April 14]. Available at: <https://www.isth.org/news/556057/ISTH-Statement-on-AstraZeneca-COVID-19-Vaccine-and-Thrombosis.htm>.
18. HSE National Immunisation Office. Clinical Guidance for COVID-19 Vaccination 2021; [internet]. [cited 2021 April 14]. Available at: <https://www.hse.ie/eng/health/immunisation/hcpinfo/covid19vaccineinfo4hps/clinicalguidance.pdf>.
19. Erskine D. Using COVID-19 vaccines in patients with anticoagulation and bleeding disorders. 2021; Specialist Pharmacy Service [internet]. [cited 2021 April 14]. Available at: <https://www.sps.nhs.uk/articles/using-covid-19-vaccines-in-patients-with-anticoagulation-and-bleeding-disorders/>.

SHPN: (ACI) 210352 TRIM: ACI/D21/695-17