

FAMILY NAME		MRN
GIVEN NAME		<input type="checkbox"/> MALE <input type="checkbox"/> FEMALE
D.O.B. ____/____/____	M.O.	
ADDRESS		
LOCATION / WARD		
COMPLETE ALL DETAILS OR AFFIX PATIENT LABEL HERE		

Facility:

DELIRIUM SCREEN FOR OLDER ADULTS

This form incorporates the **Abbreviated Mental Test scores (AMTS)**, **Delirium Risk Assessment Tool (DRAT)** and **Confusion Assessment Method (CAM)**.

Abbreviated Mental Test Score (AMTS)

Establish baseline cognition by completing the Abbreviated Mental Test OR MMSE for all presentations 65 years + (45+ ATSI). Repeat with any change in cognition behaviour or LOC. Score 1 for each correct answer.



QUESTION	Time			
	Date			
1. How old are you?				
2. What is the time (nearest hour)?				
Give the patient an address and ask them to repeat it at the end of the test eg. 42 Market St				
3. What year is it?				
4. What is the name of this place?				
5. Can the patient recognise two relevant persons? (eg. nurse / doctor or relative)				
6. What is your date of birth?				
7. When did the second World War start? (1939)				
8. Who is the current Prime Minister?				
9. Count down backwards from 20 to 1				
10. Can you remember the address I gave you?				
TOTAL SCORE				
Signature				

- A score of 7 or less indicates cognitive impairment
- All patients require a Delirium Risk Assessment using DRAT (see over page)

Does the person have a history of any recent / sudden change in behaviour, cognition, loss of consciousness or functional abilities (including falls)?

Yes - Please do CAM No - Please do DRAT

Signature: _____ Date: _____
Print full name: _____ Designation: _____

Holes Punched as per AS2828.1: 2012
BINDING MARGIN - NO WRITING

NH606735 240714

DELIRIUM SCREEN FOR OLDER ADULTS

SMR060.926



Health

FAMILY NAME

MRN

GIVEN NAME

MALE FEMALE

D.O.B. ____/____/____ M.O.

ADDRESS

DELIRIUM SCREEN FOR OLDER ADULTS

LOCATION / WARD

COMPLETE ALL DETAILS OR AFFIX PATIENT LABEL HERE

Delirium Risk Assessment Tool (DRAT)

Assessment to be completed on admission, pre & post op, and when there is a change in behaviour

Pre morbid RISK factors Tick & add score

- ≥ 70 yrs**
 - PLUS**
 - Visual impairment** (unable to read large print on newspaper with glasses)
 - Severe illness** (nurses' opinion including mental illness / depression)
 - Cognitive impairment** AMTS < 7/10 or MMSE < 25/30 or past history of memory or cognitive deficit
 - Dehydration** (scanty, concentrated urine, fever, thirst, dry mucous membranes or raised creatinine/urea)
- If your patient is ≥ 70 yrs and has at least one of the above risk factors = RISK of Delirium**

Precipitating factors

WARNING: these factors increase risk

- Mechanical restraint
- Malnutrition
- 3 new medications added in 24hrs
- IDC
- Iatrogenic event (procedure, infection complications, falls etc)

IF CHANGE IN BEHAVIOUR – RECOMMENDED INVESTIGATIONS

- | | | | | | | |
|-------------------------------------|--|--|---|---|--|-------------------------------------|
| <input type="checkbox"/> CAM | <input type="checkbox"/> Medical review | <input type="checkbox"/> History (incl. family) | <input type="checkbox"/> Physical exam | <input type="checkbox"/> Medication review | <input type="checkbox"/> Bloods | <input type="checkbox"/> MSU |
|-------------------------------------|--|--|---|---|--|-------------------------------------|

CONFUSION ASSESSMENT METHOD (CAM)

The CAM is a validated tool to be used in assisting with the differential diagnosis of delirium. It should be used for any older person who appears to be disorientated / confused or who has any change in behaviour or LOC. It is important that the CAM is used in conjunction with a formal cognitive assessment (eg AMT/MMSE), good clinical and medical assessment, together with baseline cognition information from carers/family or the community or residential aged care service.

DELIRIUM SCREENING TOOL

		No	Yes	Uncertain, Specify: _____	Is there evidence of an acute change in mental status from the patient's baseline? If so, did the abnormal behaviour fluctuate during the day?	e.g. tend to come and go or increase and decrease in severity
1	Acute onset and fluctuating course					
2	Inattention				Did the patient have difficulty focussing attention during the interview?	e.g. being easily distracted or having difficulty keeping track of what was being said
3	Disorganised thinking				Was the patient's thinking disorganised or organised?	e.g. rambling or irrelevant conversation, unclear or illogical flow of ideas, or unpredictable switching from one subject to another?
4	Altered level of consciousness				Overall, how would you rate the patient's level of consciousness?	Altered e.g. vigilant, lethargic, stupor, coma, uncertain.

Delirium is present if features 1 and 2 AND either 3 or 4 are present

Delirium symptoms: not present present Date: / /

Medical Officer notified? Yes No

Signature: _____ Date: _____

Print full name: _____ Designation: _____

Holes Punched as per AS2628.1: 2012
BINDING MARGIN - NO WRITING



240714