

Major Trauma in NSW: 2018-19

A report from the NSW Trauma Registry

August 2020

The Agency for Clinical Innovation (ACI) is the lead agency for innovation in clinical care.

We bring consumers, clinicians and healthcare managers together to support the design, assessment and implementation of clinical innovations across the NSW public health system to change the way that care is delivered.

The ACI's clinical networks, institutes and taskforces are chaired by senior clinicians and consumers who have a keen interest and track record in innovative clinical care.

We also work closely with the Ministry of Health and the four other pillars of NSW Health to pilot, scale and spread solutions to healthcare system-wide challenges. We seek to improve the care and outcomes for patients by re-designing and transforming the NSW public health system.

Our innovations are:

- person-centred
- clinically-led
- evidence-based
- value-driven.

www.aci.health.nsw.gov.au

AGENCY FOR CLINICAL INNOVATION

Level 3, 1 Reserve Road
St Leonards NSW 2065

Locked Bag 2030, St Leonards NSW 1590
T +61 2 9464 4666 | F +61 2 9464 4728
E aci-info@nsw.gov.au | aci.health.nsw.gov.au

SHPN (ACI) 200383, ISBN 978-1-76081-451-9.

Produced by: NSW Institute of Trauma and Injury Management

Further copies of this publication can be obtained from
the Agency for Clinical Innovation website at www.aci.health.nsw.gov.au

Suggested citation: NSW Agency for Clinical Innovation. Major Trauma in NSW: 2018-19. Sydney: ACI; 2020.

Disclaimer: Content within this publication was accurate at the time of publication. This work is copyright. It may be reproduced in whole or part for study or training purposes subject to the inclusion of an acknowledgment of the source. It may not be reproduced for commercial usage or sale. Reproduction for purposes other than those indicated above, requires written permission from the Agency for Clinical Innovation.

Version: 1 **Trim:** ACI/D20/2293

© Agency for Clinical Innovation 2020

Executive summary

Each year there are over 30,000 admissions for traumatic injuries in NSW, of which over 4200 were considered major trauma and admitted to a trauma service. How the NSW trauma system responds to these patients is critical for their long-term outcome and quality of life and for reducing the overall financial and social cost of trauma to individuals and the community.

A key priority for the NSW Institute of Trauma and Injury Management (ITIM), within the Agency for Clinical Innovation, is to monitor the effectiveness of the NSW trauma system response for these major trauma patients. This group of patients places the greatest demand on the trauma system and other agencies and services, not simply for healthcare, but for a wide range of needs.

This report describes how the NSW trauma system responded to major trauma patients, from the time of injury and provision of pre-hospital services, through to in-hospital services provided at a NSW trauma service. The finding from the report assists ITIM in determining whether the NSW trauma system is functioning effectively, to ensure that the right patient arrived at the right hospital in a timely matter.

The report helps us understand the nature of the injuries sustained and how they occurred. The findings are used by various agencies concerned with minimising the likelihood and effects of traumatic injury and contributing to safety and injury prevention efforts. Data from the NSW Trauma Registry is used by ITIM to provide advice and feedback to clinicians and other stakeholders and enables research into patterns of service demand and staffing.¹ This data also supports benchmarking and performance improvement activities.

It is important to note that this report does not represent all injuries in NSW, nor does it represent the full work or caseload of trauma services in hospitals or the full set of data recorded in hospital trauma registries.

2018-19 report highlights for major trauma in NSW

- 4140 major trauma patients resulted in 4252 major trauma admissions.
- Average age was 53 years old.
- Males were 2.4 times more likely to be injured than females.
- Case fatality rate for Injury Severity Score (ISS) >12 was 9.1%.
- Females had a higher case fatality rate for ISS > 12 (11.0%), compared to males (8.3%).
- Falls accounted for 44.0% of injuries, exceeding transport incidents (37.4%).
- The most common serious injury was 'three or more fractured ribs without flail' (24.2%).
- 27.0% of major traumas were sustained in a rural area.
- The greatest proportion of traumatic injuries were caused by falls in the metro setting (50.1%) and by transport incidents in the rural setting (49.1%).
- Pedestrian traumas had a significantly higher case fatality rate (18.9%) than all other forms of road trauma.
- 58.3% of major trauma patients sustained injuries to the head or neck body region, and 51.5% to the chest region.
- Overall data completeness of mandatory elements was 95.0%.

Contents

Executive summary	iii
Monitoring the NSW trauma system.....	5
NSW trauma system	5
Mandate for trauma data	5
NSW Trauma Registry.....	5
NSW trauma services.....	6
Methodology	7
Inclusion criteria	7
Exclusion criteria	7
Data quality	8
Data privacy – reporting of small numbers	8
Metropolitan and rural categorisation	8
The Injury Severity Score and Abbreviated Injury Score	9
Revised Trauma Score.....	9
Major trauma patients.....	10
Introduction.....	10
Summary profile	10
Age and sex	12
Mechanism of injury	16
Transport incidents by place of occurrence	24
Time and day of injury	26
Injuries.....	27
Injury Severity Score	29
Pre-hospital time	31
Mode of transport	32
Major trauma admissions.....	34
Introduction.....	34
Overall burden of trauma.....	34
Facility overview	35
Admission type	36
Revised Trauma Score.....	36
Trauma team activation	37

Vital signs on arrival to the emergency department.....	39
Intensive care unit admissions	41
Length of stay.....	42
Procedures	44
Discharge destination of survivors.....	45
References	47
Glossary	48
Abbreviations.....	50
Acknowledgements	51
Appendices	52
Appendix 1: Data completeness.....	53
Appendix 2: Adult major trauma service summaries	57
Appendix 3: Paediatric major trauma service summaries.....	71
Appendix 4: Regional trauma service summaries	77
Appendix 5: Calculation of the Injury Severity Score.....	97
Appendix 6: Australian Statistical Geography Standard Remoteness Areas.....	98

List of tables

Table 1: Record of data exclusions	7
Table 2: Revised Trauma Score points system.....	9
Table 3: Summary statistics for major trauma and mortality	10
Table 4: Major trauma patients, injury and mortality by age group	12
Table 5: Type of injury	16
Table 6: Mechanism of injury by age.....	17
Table 7: Mechanism of injury by location	20
Table 8: Falls in detail	21
Table 9: Transport incidents in detail.....	22
Table 10: Assaults in detail	22
Table 11: All other mechanisms in detail.....	23
Table 12: Transport incidents by location of injury.....	24
Table 13: Road trauma in detail	24
Table 14: Other transport incidents in detail.....	25
Table 15: Top 5 injuries with an AIS severity >2.....	27
Table 16: Number of ISS body regions injured with an AIS severity >2.....	28
Table 17: Single body region versus polytrauma with an AIS severity >2.....	28
Table 18: Major trauma patients by ISS group	29

Table 19: Median time of injury to arrival at a designated trauma service	31
Table 20: Median time of injury to arrival at definitive care	31
Table 21: Mode of transport to definitive care	32
Table 22: Overview of trauma service admissions	35
Table 23: Number of admissions by type	36
Table 24: Overview of ICU and hospital length of stay	42
Table 25: ICU and hospital length of stay by ISS	42
Table 26: ICU and hospital length of stay by age	43
Table 27: Procedures performed by type	44
Table 28: Detailed data completeness by facility	54
Table 29: Trauma data profile, John Hunter Hospital	57
Table 30: Trauma data profile, Liverpool Hospital	59
Table 31: Trauma data profile, Royal North Shore Hospital	61
Table 32: Trauma data profile, Royal Prince Alfred Hospital	63
Table 33: Trauma data profile, St George Hospital	65
Table 34: Trauma data profile, St Vincent's Hospital	67
Table 35: Trauma data profile, Westmead Hospital	69
Table 36: Trauma data profile, John Hunter Children's Hospital	71
Table 37: Trauma data profile, Sydney Children's Hospital	73
Table 38: Trauma data profile, The Children's Hospital at Westmead	75
Table 39: Trauma data profile, Coffs Harbour Base Hospital	77
Table 40: Trauma data profile, Gosford Hospital	79
Table 41: Trauma data profile, Lismore Base Hospital	81
Table 42: Trauma data profile, Nepean Hospital	83
Table 43: Trauma data profile, Orange Base Hospital	85
Table 44: Trauma data profile, Port Macquarie Base Hospital	87
Table 45: Trauma data profile, Tamworth Base Hospital	89
Table 46: Trauma data profile, The Tweed Hospital	91
Table 47: Trauma data profile, Wagga Wagga Base Hospital	93
Table 48: Trauma data profile, Wollongong Hospital	95
Table 49: ISS calculation example	97

List of figures

Figure 1: Five-year trend in the number of major trauma patients	11
Figure 2: Five-year trend in the number of major trauma patients by facility type	11
Figure 3: Number of major trauma patients by age, sex and mortality	13
Figure 4: Age-specific injury rate by age and sex	13
Figure 5: Case fatality rate by age and sex (ISS >12)	14
Figure 6: Age-specific mortality rate by age and sex	14
Figure 7: Five-year trend of case fatality rate by facility type (ISS >12)	15
Figure 8: Mechanism of injury	16
Figure 9: Five-year trend in the number of falls and transport incidents	18
Figure 10: Mechanism of injury by age	18
Figure 11: Mechanism of injury as a percentage by age	19
Figure 12: Age-specific injury rate by mechanism of injury	19
Figure 13: Mechanism of injury by location	20
Figure 14: Mechanism of injury as a percentage by location	21
Figure 15: Number of patients by time of injury	26
Figure 16: Number of patients by day of injury	26
Figure 17: All injuries by ISS body region	27
Figure 18: Number of major trauma patients by ISS group and sex	29
Figure 19: Case fatality rate by ISS and sex	30
Figure 20: Mode of transport to definitive care when transported direct from the scene of injury, by injury location	32
Figure 21: Mode of transport to definitive care when transferred from another acute care facility, by injury location	33
Figure 22: Number of patients with an estimated ISS >5 by local health district	34
Figure 23: Average Revised Trauma Score by ISS group	36
Figure 24: Trauma team activation by admission type	37
Figure 25: Trauma team activation by ISS group	38
Figure 26: Initial systolic blood pressure on arrival to the ED and mortality	39
Figure 27: Initial Glasgow Coma Scale on arrival to the ED and mortality	39
Figure 28: Initial temperature on arrival to ED and mortality	40
Figure 29: ICU admission by ISS group	41
Figure 30: Procedures performed by ISS group	44
Figure 31: Five-year trend of procedures performed	45
Figure 32: Discharge destination of survivors	45
Figure 33: Discharge destination of survivors by ISS group	46
Figure 34: Overall data completeness by facility	53
Figure 35: Map of 2016 Remoteness areas for Australia	98

Monitoring the NSW trauma system

NSW trauma system

The primary function of the NSW trauma system is to facilitate and coordinate an organised multidisciplinary system response with the aim of reducing the burden of injury on patients, their families and the broader health system. The trauma system encompasses a continuum of care that provides traumatically injured patients with the greatest likelihood of returning to their pre-injury level of function within the community.

This continuum of care includes injury prevention, pre-hospital coordination and care, appropriate triage and transport, emergency department trauma care, trauma service team activation, surgical intervention, intensive, critical and general in-hospital care, rehabilitation services, allied health and medical care follow up.

The overall goal of the NSW trauma system is to decrease the incidence and severity of injury and to ensure optimal, accessible and equitable care to improve health outcomes for those who are injured. The main objective of the trauma system is to get the 'right patient to the right hospital at the right time, receiving the right care'. To meet this objective, designated trauma services need to have appropriate resources to meet the complex needs of the injured patient.

Mandate for trauma data

The *NSW Trauma Services Plan* outlines the role of ITIM, including monitoring and reporting on the performance of individual trauma services to ensure that performance is consistent with the standard of care and to manage a statewide clinical injury data collection process.²

The plan also positions ITIM to develop partnerships with injury stakeholders, such as the local health districts, NSW Ambulance, State Insurance Regulatory Authority, State Coroner, Clinical Excellence Commission and universities. These partnerships aim to build an improved critical mass for research and education across the spectrum of trauma prevention care and rehabilitation. The collection of trauma data is an important aspect of these activities.

To enable ITIM to achieve its monitoring and reporting role, NSW trauma services are required to contribute data as outlined in *Improving the quality of trauma care in NSW: trauma services model of care*.³

NSW Trauma Registry

ITIM is responsible for managing the collection of data about moderate to critically injured people admitted to trauma services in NSW. Data collected is held securely in the NSW Trauma Registry.¹ Data in the registry is submitted from each of the designated NSW trauma services.

The NSW Trauma Registry contains de-identified patient records but does not hold data for every injured person admitted to hospital in NSW. Data is only included for patients with the greatest needs, the most seriously injured, who are treated at a designated NSW trauma service. This data is known as the NSW Trauma Minimum Data Set and forms the basis of data analysis and reporting activities at ITIM. As the scope of the current data collection is restricted to these designated hospitals, there may be some data for trauma admissions to other hospitals that are not included in the NSW Trauma Registry.

NSW trauma services

The NSW trauma system consists of seven adult major trauma services, three paediatric major trauma services and ten regional trauma services. All of the services contributed data to the NSW Trauma Registry used in this report.

Adult major trauma services

- John Hunter Hospital, Hunter New England Local Health District
- Liverpool Hospital, South Western Sydney Local Health District
- Royal North Shore Hospital, Northern Sydney Local Health District
- Royal Prince Alfred Hospital, Sydney Local Health District
- St George Hospital, South Eastern Sydney Local Health District
- St Vincent's Hospital, St Vincent's Health Network
- Westmead Hospital, Western Sydney Local Health District

Paediatric major trauma services

- John Hunter Children's Hospital, Hunter New England Local Health District
- Sydney Children's Hospital, Sydney Children's Hospital Network
- The Children's Hospital at Westmead, Sydney Children's Hospital Network

Regional trauma services

- Coffs Harbour Health Campus, Mid North Coast Local Health District
- Gosford Hospital, Central Coast Local Health District
- Lismore Base Hospital, Northern NSW Local Health District
- Nepean Hospital, Nepean Blue Mountains Local Health District
- Orange Health Service, Western NSW Local Health District
- Port Macquarie Base Hospital, Mid North Coast Local Health District
- Tamworth Hospital, Hunter New England Local Health District
- The Tweed Hospital, Northern NSW Local Health District
- Wagga Wagga Rural Referral Hospital, Murrumbidgee Local Health District
- Wollongong Hospital, Illawarra Shoalhaven Local Health District

Methodology

This report is compiled from data submitted by the reporting facilities to the NSW Trauma Registry in accordance with the inclusion and exclusion criteria detailed below. Data for this report was extracted from the NSW Trauma Registry on 6 May 2020.

Inclusion criteria

All major trauma patient records from the NSW Trauma Registry, where the date of injury occurred between 1 July 2018 and 30 June 2019, are included in this report.

Major trauma is defined as all patients of any age, who were admitted to a NSW trauma service within seven days of sustaining an injury, and who:

- had an Injury Severity Score (ISS) >12 (moderate to critically injured), or
- were admitted to an intensive care unit (irrespective of ISS) following injury, or
- died in hospital (irrespective of ISS) following injury.

As a result of these criteria, patient records submitted for inclusion in this report do not represent all injuries in NSW, nor do they represent the full work or caseload of trauma services in hospitals, nor the full set of data recorded in hospital trauma registries.

Exclusion criteria

The criteria for excluding a patient record from this report are as follows.

- Patients not admitted to a designated NSW trauma service.
- Patients admitted to a designated NSW trauma service greater than seven days after sustaining an injury.
- Patients who die with an isolated fractured neck of femur injury sustained from a fall from a standing height (<1 metre).*
- Patients aged 65 years or older who die with minor soft tissue injury only.†

Records have also been excluded from this report if the *Outcome (survived or died)* data element is missing or invalid data recorded in the registry.

Based on the criteria outlined above, a total of 49 records were excluded as outlined in Table 1.

Table 1: Record of data exclusions

Data criteria	Excluded	Remaining records
Data extracted (6 May 2020)	N/A	4301
Date of admission >7 days from injury	42	4259
Isolated neck of femur injuries (with outcome = died)	4	4255
Over 65yrs old and died with minor soft tissue injury only	1	4254
Missing outcome	2	4252
Total remaining records		4252

* See [Glossary](#) for definition of an isolated fractured neck of femur injury.

† See [Glossary](#) for definition of a minor soft tissue injury.

Data quality

Data submitted to the NSW Trauma Registry is subject to rigorous checking and validation by ITIM and the reporting facilities while missing or invalid data is flagged and returned to individual trauma services for completion and validation.

The average data completeness by the trauma facilities was 95.0%, an increase of 4.9% from last year. A detailed breakdown of data completeness by trauma facility is available in [Appendix 1](#).

Data privacy – reporting of small numbers

The Australian Statistical Information Management Committee guidelines suggest that statistical results involving small numbers (<5) can be presented if the population from which they are drawn is more than 1000 people, as the likelihood of identifying an individual would be very small.⁴ This guideline is used for data reporting of small numbers by Health Stats NSW and the Agency for Clinical Innovation.⁵

For the purpose of this report, the demographic population is identified as either:

- those potentially injured within NSW (millions) when reporting on trauma patients as a whole; or
- those potentially injured within a facility's catchment area (thousands) when reporting on admissions to a trauma facility, such as in the facility summaries at Appendix 2-4.

The risk of a breach of data privacy is further mitigated by not aligning demographic information where the numbers are small (such as age groups in a regional area) with other potentially identifiable data fields such as the mechanism of injury or outcome.

Metropolitan and rural categorisation

Various data elements within the report are categorised as either 'metropolitan' or 'rural'. These categories are derived using the postcode of injury and the Australian Statistical Geography Standard (ASGS) Remoteness Areas (RA).⁶ The ASGS-RA is based on the *Accessibility and Remoteness Index of Australia* which defines locations in terms of remoteness, i.e. the physical distance of a location from the nearest urban centre (access to goods and services) based on population size.⁷

The ASGS-RA consists of five categories:

- major cities
- inner regional
- outer regional
- remote
- very remote.

For the purpose of this report, all locations with the ASGS-RA classification of 'major cities' are listed as 'metropolitan'. All other ASGS-RA classified locations are combined and listed as 'rural'. See [Appendix 6](#) for further information on the ASGS-RA categorisation of NSW.

The Injury Severity Score and Abbreviated Injury Score

One of the key criteria for inclusion in this report is an Injury Severity Score (ISS) >12. The ISS is an internationally recognised scoring system which correlates with mortality, morbidity and other measures of severity. The ISS is calculated based on an anatomical injury severity classification, the Abbreviated Injury Scale (AIS). The AIS classifies individual injuries by body region on a six-point severity scale from minor (1) to maximum (6 - currently untreatable injury). The NSW Trauma Registry uses the AIS 2005 (Update 2008) dictionary.⁸

The AIS is used by accredited staff at each hospital to score individual patient injuries and their severity. It provides a common tool for comparing and selecting patient records for inclusion in the NSW Trauma Registry. Scoring is undertaken retrospectively but usually within 24-48 hours after admission to allow for the identification of all injuries. On initial evaluation, these patients typically have abnormal vital signs or a significant anatomical injury.

Injuries are individually allocated to one of six body regions, and the severities of the top three injuries in different body regions are used to calculate the ISS. The ISS, along with the body regions and injury and severity codes, used in this calculation, are recorded in the NSW Trauma Registry (see [Appendix 5](#) for further detail).

The calculated ISS value ranges from 1-75. Serious to critically injured trauma patients are defined as those patients with an ISS >15, which is an internationally recognised indicator of serious injury.

In this report, the ISS is reported in ranges:

- 13-15 (moderate injury)
- 16-24 (serious injury)
- 25-40 (severe injury)
- 41-75 (critical injury).

Revised Trauma Score

The Revised Trauma Score is a physiological scoring system used as a predictor of mortality in trauma populations. It consists of data from the first set of vital signs obtained on arrival at the hospital, including the Glasgow Coma Scale, systolic blood pressure and respiratory rate. Each element is scored with a weighting as outlined in Table 2.⁹ Values for the Revised Trauma Score are in the range 0 to 7.8408. The lower the score, the higher the likelihood of death.

Table 2: Revised Trauma Score points system

Glasgow Coma Scale	Systolic blood pressure	Respiratory rate	Points
15-13	>89	10-29	4
12-9	76-89	>29	3
8-6	50-75	6-9	2
5-4	1-49	1-5	1
3	0	0	0

Revised Trauma Score = 0.9368 Glasgow Coma Scale + 0.7326 systolic blood pressure + 0.2908 respiratory rate

Major trauma patients

Introduction

The information in this section of the report is based on the number of major trauma patients who received care in a NSW trauma service (n=4140), not the number of major trauma admissions (n=4252), as some patients were treated in more than one NSW reporting facility. Trauma admission data (hospital activity) is discussed in detail in the [Major trauma admissions](#) section.

Summary profile

During the period 1 July 2018 to 30 June 2019, there were 4140 major trauma patients treated at NSW trauma services. Of these, 73% (n=2824) were injured in a metropolitan location, and 316 died (case fatality rate for ISS >12 of 9.1%). The age-standardised injury rate was 48.9 per 100,000 persons, and the age-standardised death rate was 4.2 per 100,000 persons for all ISS.* The standardised mortality ratio† was 3.8, indicating that the proportion of deaths in major trauma patients during the reporting period was more than four times greater than that of the general Australian population (Table 3).

Table 3: Summary statistics for major trauma and mortality


Summary statistics	Value
Total number of patients injured overall	4140
Total number of patients injured with ISS >12	3464
Injury rate per 100,000 persons (age-standardised)	47.7 (95% CI 46.2 - 49.2)
Location of injury (metropolitan / rural)	2824 (73%) / 1045 (27%)
Number of male / female patients	2934 (70.9%) / 1206 (29.1%)
Total number of deaths overall	406 (9.8%)
Total number of traumatic deaths on arrival	21 (0.5%)
Total number of deaths with ISS >12 (case fatality rate)	316 (9.1%)
Total number of deaths with ISS >12 excluding traumatic deaths on arrival	299 (8.7%)
Death rate per 100,000 persons (age-standardised) all ISS	4.2 (95% CI 3.8 - 4.6)
Standardised mortality ratio (SMR)	3.8 (95% CI 3.4 - 4.2)
Average age (years)	52.8 (95% CI 52 - 53.6)
Average ISS (all ISS / ISS >12)	18.4 / 20.5

* See [Glossary](#) for definition of age-standardised rate. Annualised rate given as per 100,000 persons, standardised to the Australian population at 30 June 2001.¹⁰

† See [Glossary](#) for definition of the standardised mortality ratio. Standardised to the Australian population at 30 June 2001.¹⁰


From 2014-15 to 2018-19 financial years, there has been a steady minor increase (6.0%) in the overall number of major trauma patients, except for 2016-17 (Figure 1).

Figure 1: Five-year trend in the number of major trauma patients (n=20,005)



The largest proportional increase from 2017-18 compared to 2018-19 was seen in the paediatric trauma services (9.0%), compared to regional trauma services (5.2%) and adult major trauma services (0%), as seen in Figure 2.

Figure 2: Five-year trend in the number of major trauma patients by facility type (n=20,005)



Age and sex

The average age of a major trauma patient in NSW during the reporting period was 52.8 years old. Table 4 outlines the age distribution, which demonstrates a sharp rise in the incidence of major trauma in the geriatric population,* especially those aged 75 years and older, as it has in previous years. The 75 years and older age group also had a higher case fatality rate for ISS >12 (20%, n=827) compared with the under 75 years age group (5.7%, n=2637).


Table 4: Major trauma patients, injury and mortality by age group (n=4140)

Age group (years)	Number of injured (% of total)	Cumulative number of injured (% of total)	Age-specific injury rate per 100,000	Age-specific death rate per 100,000	Case fatality rate (ISS >12)
0-4	96 (2.3%)	96 (2.3%)	19.3	2.2	12.2%
5-9	48 (1.2%)	144 (3.5%)	9.5	0.2	0%
10-14	78 (1.9%)	222 (5.4%)	16.7	0.4	3.3%
15-19	196 (4.7%)	418 (10.1%)	41.8	2.3	5.5%
20-24	285 (6.9%)	703 (17.0%)	52.7	3.5	5.9%
25-29	235 (5.7%)	938 (22.7%)	39.7	2.4	7.0%
30-34	202 (4.9%)	1140 (27.5%)	34.6	0.3	1.3%
35-39	217 (5.2%)	1357 (32.8%)	40.7	2.3	5.6%
40-44	214 (5.2%)	1571 (37.9%)	42.0	1.4	3.7%
45-49	254 (6.1%)	1825 (44.1%)	49.4	1.9	4.0%
50-54	257 (6.2%)	2082 (50.3%)	52.9	1.9	3.9%
55-59	261 (6.3%)	2343 (56.6%)	53.8	2.3	4.0%
60-64	286 (6.9%)	2629 (63.5%)	66.3	3.7	5.6%
65-69	233 (5.6%)	2862 (69.1%)	60.6	6.0	9.5%
70-74	259 (6.3%)	3121 (75.4%)	82.4	9.5	11.2%
75-79	257 (6.2%)	3378 (81.6%)	114.8	18.8	16.2%
80-84	310 (7.5%)	3688 (89.1%)	196.0	37.9	17.7%
85+	452 (10.9%)	4140 (100%)	266.7	74.4	23.5%

Males have a higher number of major trauma injuries (n=2934) compared to females (n=1206), except in the 85 years and over age groups (Figure 3). Overall, males are 2.4 times more likely to be injured than females.


* Geriatric defined as aged 65 years and older.

Figure 3: Number of major trauma patients by age, sex and mortality (n=4140)



The age-specific injury rate* for males ranged from 4.9 to 134.1 per 100,000 persons, and in females ranged 3.72 to 126.1 per 100,000 persons (Figure 4).

Figure 4: Age-specific injury rate by age and sex (n=4140)



* See [Glossary](#) for definition of the age-specific rate, based on the estimated NSW population during the reporting period (end of December).¹¹

The case fatality rate for ISS >12 (n=3464) was higher for females (11.0%, n=111) than for males (8.3%, n=205), with the overall case fatality rate being 9.1% (Figure 5).

Figure 5: Case fatality rate by age and sex (ISS >12) (n=3464)



While females have a higher case fatality rate for ISS >12, males have a higher age-specific mortality rate per 100,000 persons (Figure 6).

Figure 6: Age-specific mortality rate by age and sex (n=4140)





Figure 7 shows the five-year trend of case fatality rates (ISS >12) for the three facility types. Overall, the case fatality rate has decreased from 10.5% to 9.1% during the last five years.

Figure 7: Five-year trend of case fatality rate by facility type (ISS >12) (n=17,408)



Mechanism of injury

The vast majority of major trauma in NSW in the reporting period was caused by blunt injuries (94.5%), such as falls and transport incidents (Table 5).

Table 5: Type of injury (n=4050)*


Type of injury	Number of patients (% of total)	Case fatality rate (ISS >12)
Blunt	3826 (94.5%)	8.85%
Penetrating	224 (5.5%)	8.93%

The top three mechanisms of major trauma were:

- falls (44.0%, n=1820)
- transport incidents (37.3%, n=1547) out of which 1184 were road trauma incidents[†]
- assaults (6.4%, n=264).

All other mechanisms combined accounted for 12.3% (n=509) (Figure 8). The distribution of these mechanisms of injury by age group is demonstrated in Table 6.

Figure 8: Mechanism of injury (n=4140)



* Only blunt and penetrating injury types are reported. Other injury types are recorded as 'N/A' or 'Unknown'.


† See [Glossary](#) for the definition of road trauma.

Table 6: Mechanism of injury by age (n=4140)

Age group	Number of patients (age-specific rate per 100,000)			
	Falls	Transport incident	Assault	All other mechanisms
0-4	33 (6.6)	17 (3.4)	12 (2.4)	34 (6.8)
5-9	18 (3.5)	16 (3.1)	1 (0.2)	13 (2.5)
10-14	18 (3.7)	45 (9.3)	0 (0)	15 (3.1)
15-19	31 (6.5)	120 (25.3)	14 (3)	31 (6.5)
20-24	48 (8.7)	152 (27.4)	36 (6.5)	49 (8.8)
25-29	39 (6.4)	120 (19.7)	28 (4.6)	48 (7.9)
30-34	25 (4.2)	96 (16.1)	38 (6.4)	43 (7.2)
35-39	40 (7.1)	102 (18.2)	34 (6.1)	41 (7.3)
40-44	59 (11.6)	96 (18.9)	23 (4.5)	36 (7.1)
45-49	75 (14.3)	118 (22.5)	20 (3.8)	41 (7.8)
50-54	81 (16.9)	113 (23.6)	21 (4.4)	42 (8.8)
55-59	105 (21.3)	105 (21.3)	12 (2.4)	39 (7.9)
60-64	124 (28)	131 (29.5)	12 (2.7)	19 (4.3)
65-69	134 (34.3)	83 (21.2)	5 (1.3)	11 (2.8)
70-74	184 (54.6)	59 (17.5)	2 (0.6)	14 (4.2)
75-79	173 (73.4)	69 (29.3)	2 (0.8)	13 (5.5)
80-84	240 (144.9)	58 (35)	3 (1.8)	9 (5.4)
85+	393 (226.3)	47 (27.1)	1 (0.6)	11 (6.3)
Total	1820	1547	264	509

Figure 9 shows the overall five-year trend of increasing falls and the decreasing transport incidents resulting in major trauma, although there is variability from year to year.

Figure 9: Five-year trend in the number of falls and transport incidents (n=20,005)



The greatest burden of major trauma for people aged 65 years and older is falls (74.4%, n=1124), whilst for those aged under 65 years it is transport incidents (46.8% n=1231).

The highest incidence of assaults is in the 30-34 years age group (n=38) closely followed by the 20-24 years age group (n=36). See Table 6 and Figures 10-12.

Figure 10: Mechanism of injury by age (n=4,140)




Figure 11: Mechanism of injury as a percentage by age (n=4140)





Figure 12: Age-specific injury rate by mechanism of injury (n=4140)



Falls were responsible for 50.1% of the injuries in the metropolitan area, compared to 31.0% in rural areas. Transport incidents accounted for a higher percentage of injuries in rural areas, 49.1%, than in metropolitan areas, 32.0% (Table 7, Figures 13-14).

Table 7: Mechanism of injury by location (n=3869)

Mechanism of injury	Metropolitan (% of metropolitan)	Rural (% of rural)
Falls	1415 (50.1%)	324 (31.0%)
Transport Incident	904 (32.0%)	513 (49.1%)
Assault	176 (6.2%)	61 (5.8%)
All other mechanisms	329 (11.7%)	147 (14.1%)
Total	2824 (73.0%)	1045 (27.0%)

Figure 13: Mechanism of injury by location (n=3869)





Figure 14: Mechanism of injury as a percentage by location (n=3869)



The mechanisms of injury are outlined in more detail in Tables 8-11.

Falls from a standing height (lowest level, <1m) accounted for the greatest burden of all heights. They accounted for 61.3% of all falls, 26.1% of all trauma mechanisms, 30.5% of all trauma deaths, and had case fatality rate for ISS >12 of 13.8% (Table 8).

Table 8: Falls in detail (n=1820)

Mechanism	Number injured (% of all mechanisms)	Number of deaths, ISS >12 (case fatality rate)	Percentage of all trauma deaths
Low fall (<1m)	1116 (27%)	124 (13.8%)	30.5%
Medium fall (1-5m)	561 (13.6%)	37 (7.3%)	9.1%
High fall (>5m)	74 (1.8%)	4 (5.8%)	1.0%
Unspecified fall	69 (1.7%)	11 (25.0%)	2.7%
Total	1820 (44.0%)	176 (11.6%)	43.3%

Pedestrian trauma had the highest case fatality rate for ISS >12 (14.1%) (Table 9) well above other forms of transport incidents. Pedestrian mortality rate was 6.7% of all the trauma deaths

Table 9: Transport incidents in detail (n=1547)

Mechanism	Number injured (% of all mechanisms)	Number of deaths, ISS >12 (case fatality rate)	Percentage of all trauma deaths
Car occupant	576 (13.9%)	39 (8.1%)	9.6%
Motorcycle rider	414 (10.0%)	6 (1.6%)	1.5%
Pedal cyclist	211 (5.1%)	5 (2.6%)	1.2%
Pedestrian	210 (5.1%)	27 (14.1%)	6.7%
All other transport	136 (3.3%)	3 (2.5%)	0.7%
Total	1547 (37.4%)	80 (5.9%)	19.7%

The most common mechanisms of injury in the assault group were assaults involving bodily force (n=96, 2.3% of all mechanisms), assault by knife (n=72, 1.7%), and assault by blunt object (n=33, 0.8%). Notably, assault by knife increased by 38% from last year. Assault by firearm remains low with only nine recorded cases (0.2%) (Table 10).

Table 10: Assaults in detail (n=264)

Mechanism	Number injured (% of all mechanisms)	Number of deaths, ISS >12 (case fatality rate)	Percentage of all trauma deaths
Assault by bodily force	96 (2.3%)	4 (5.3%)	1.0%
Assault by knife	72 (1.7%)	6 (11.1%)	1.5%
All other assaults	54 (1.3%)	4 (10.0%)	1.0%
Assault by blunt object	33 (0.8%)	0 (0%)	0%
Assault by firearm	9 (0.2%)	1 (12.5%)	0.2%
Total	264 (6.4%)	15 (7.3%)	3.7%

Of the other mechanisms of injury, self-harm was the most common (n=173, 4.2%). Drownings and 'Other' had the highest case fatality rate for ISS >12 in the 'all other mechanisms' group at 35.7% and 22.2% respectively (Table 11).

Table 11: All other mechanisms in detail (n=509)

Mechanism	Number injured (% of all mechanisms)	Number of deaths, ISS >12 (case fatality rate)	Percentage of all trauma deaths
Self-harm	173 (4.2%)	20 (15.9%)	4.9%
Burns	50 (1.2%)	0 (0%)	0%
Drownings	37 (0.9%)	10 (35.7%)	2.5%
Other	53 (1.3%)	8 (22.2%)	2.0%
Animate mechanical forces	71 (1.7%)	1 (1.6%)	0.2%
Inanimate mechanical forces	125 (3.0%)	6 (6.0%)	1.5%
Total	509 (12.3%)	45 (11.1%)	11.1%

Transport incidents by place of occurrence

In order to provide a greater analysis of the burden road trauma has on the health system, a 'place of occurrence' code qualifier was applied to all transport incidents resulting in two categories: road trauma; and other transport incidents. For the purpose of this report, road trauma is defined as a transport incident which occurred on a street, highway or other paved roadway.*

Road trauma accounts for 28.6 % (n=1184) of all mechanisms of injury, the second highest behind falls at 44.0% (n=1820). The rate of 'road trauma' and 'other transport incidents' were higher in rural areas, 34.8% and 14.3% respectively, than in metropolitan areas, 25.9% and 6.1% respectively (Table 12).

Table 12: Transport incidents by location of injury (n=1547)

Mechanism of injury	Metropolitan (% of metropolitan)	Rural (% of rural)	Unknown location (% of unknown)
Road trauma	731 (25.9%)	364 (34.8%)	89 (32.8%)
Other transport incidents	173 (6.1%)	149 (14.3%)	41 (15.1%)
Total	904	513	130

Injuries to car occupants remains the highest mechanism of injury in the road trauma group (n=547, 13.2%) with pedestrian trauma having the highest case fatality rate for ISS >12 (15.4%), well above other forms of road trauma (Table 13).

Table 13: Road trauma in detail (n=1184)

Mechanism	Number injured (% of all mechanisms)	Number of deaths, ISS >12 (case fatality rate)	Percentage of all trauma deaths
Car occupant	547 (13.2%)	33 (7.2%)	8.1%
Motorcycle rider	283 (6.8%)	6 (2.3%)	1.5%
Pedestrian	176 (4.3%)	25 (15.4%)	6.2%
Pedal cyclist	146 (3.5%)	5 (3.7%)	1.2%
All other road transport	32 (0.8%)	2 (6.7%)	0.5%
Total	1184 (28.6%)	71 (6.8%)	17.5%

* See [Glossary](#) for a detailed definition of 'road trauma' and 'other transport incident'.

The most common mechanisms in the 'other transport incidents' group were motorcycle riders (n=131), other land transport incidents (n=83), pedal cyclists (n=65), and pedestrians (n=34). Of the 'other land transport' incident group, animal-rider or animal-drawn vehicle were the most common (n=51) followed by all-terrain vehicle (including quad bike) incidents (n=24) (Table 14).

Other transport incidents include incidents being reported as occurring NOT on a street, highway or other paved roadway.


Table 14: Other transport incidents in detail (n=363)

Mechanism	Number injured (% of all mechanisms)	Number of deaths, ISS >12 (case fatality rate)	Percentage of all trauma deaths
Motorcycle rider	131 (3.2%)	0 (0%)	0%
Other land transport accidents	83 (2.0%)	1 (1.4%)	0.2%
Pedal cyclist	65 (1.6%)	0 (0%)	0%
Pedestrian	34 (0.8%)	2 (6.7%)	0.5%
Car occupant	29 (0.7%)	6 (25.0%)	1.5%
Water transport accidents	10 (0.2%)	0 (0%)	0%
Air and space transport accidents	6 (0.1%)	0 (0%)	0%
Occupant of pick-up truck or van	3 (0.1%)	0 (0%)	0%
Occupant of heavy transport vehicle	2 (0%)	0 (0%)	0%
Total	363 (8.8%)	9 (2.8%)	2.2%

Time and day of injury


Of note is that 55.3% of patients (n=1888) were injured between 11am and 8pm, which resulted in peak activity in the hospitals during the afternoon and evening (Figure 15).

Figure 15: Number of patients by time of injury (n=3417)



The mean daily number of persons injured on a weekday was 10.4. The number of persons injured on weekends was higher, with a mean of 13.7 per day (Figure 16).*

Figure 16: Number of patients by day of injury (n=4140)



* In the reporting period there were 260 weekdays and 105 weekend days.

Injuries


Three or more fractured ribs without a flail segment* continued to be the most common single serious injury sustained (24.0%, n=995) where the AIS severity was greater than two (Table 15). Cerebral haematomas occupied three of the top five injuries.

Table 15: Top five injuries with an AIS severity >2 (n=4140)

Injury description (as per AIS dictionary)	AIS severity	Number of patients (% of total)
Fractured ≥ 3 ribs without flail, not further specified	3	995 (24.0%)
Cerebrum hematoma - subdural - small; moderate	4	389 (9.4%)
Cerebrum hematoma - subdural - large; massive; extensive	5	216 (5.2%)
Cerebrum hematoma - subdural - tiny	3	206 (5.0%)
Hemopneumothorax, not further specified	3	198 (4.8%)

In addition, 58.3% of major trauma patients sustained injuries to the head or neck body region, with the chest region injured in 51.5% of major trauma patients (Figure 17).

Figure 17: All injuries by ISS body region (n=4140)



* Flail is defined as three or more ribs fractured in more than one location and/or resulting in paradoxical chest movement.⁸

The majority of major trauma patients (72.7%) sustained serious injury (AIS severity >2) to only one ISS body region (Table 16).

Table 16: Number of ISS body regions injured with an AIS severity >2 (n=4140)

Number of ISS body regions injured with an AIS severity >2	Number of patients - All ISS (% of total)	Number of patients (% of total) (ISS >12)
0	325 (7.9%)	0 (0%)
1	3009 (72.7%)	2660 (76.8%)
2	680 (16.4%)	678 (19.6%)
3	112 (2.7%)	112 (3.2%)
4	12 (0.3%)	12 (0.3%)
5	2 (0%)	2 (0.1%)

21.1% of major trauma patients sustained polytrauma, defined as sustaining serious injury (AIS severity >2) in two or more ISS body regions (Table 17).¹²

Table 17: Single body region versus polytrauma with an AIS severity >2 (n=3815)

Body region	Number of patients - all ISS (% of total)	Case fatality rate (ISS >12)
Single body region	3009 (78.9%)	8.1%
Polytrauma	806 (21.1%)	12.4%

Injury Severity Score


An ISS of greater than 12 is a key identifier of a major trauma patient. The ISS correlates with mortality – the higher the ISS, the higher the mortality rate (Table 18). It is important to note that ISS is only calculated on injuries sustained and does not include other potential contributors to mortality and morbidity, such as patient's age and comorbidities. The average ISS for all major trauma patients was 18.4. This increased to 20.5 when excluding those with an ISS less than 13.

Table 18: Major trauma patients by ISS group (n=4131)

ISS group	Number of patients (% of total)	Number of deaths (case fatality rate)
ISS <13	667 (16.1%)	83 (12.4%)
ISS 13-15	957 (23.2%)	19 (2.0%)
ISS 16-24	1535 (37.2%)	60 (3.9%)
ISS 25-40	856 (20.7%)	191 (22.3%)
ISS 41-75	116 (2.8%)	46 (39.7%)


The serious injury category (ISS 16-24) contained the highest number of injured at 1535 (37.2%), followed by the moderate injury category (ISS 13-15) which had 957 (23.2%) injured and the severe injury category (ISS 25-40) had 856 (20.7%) (Figure 18).

Figure 18: Number of major trauma patients by ISS group and sex (n=4131)



Females had a higher case fatality rate than males in all the ISS groups (Figure 19). Overall the case fatality rates were higher in females than males (12.4% versus 8.8%) for all ISS.

Figure 19: Case fatality rate by ISS and sex (n=3464)



Pre-hospital time

The time from injury to arrival at a designated trauma service can have a significant impact on morbidity and mortality in the major trauma patient cohort. The regional population and geography of NSW are vastly spread, and this impacts on the variation in the time of arrival to a designated trauma service. Patients who were injured in a metropolitan region arrived at a designated trauma service faster (77 mins) than those injured in a rural location (137 mins) (Table 19).

See the [Methodology](#) section of this report for more information regarding the definitions of metropolitan and rural.

Table 19: Median time of injury to arrival at a designated trauma service (n=3891)

Location of injury	Direct from scene	Transferred from another hospital	Overall
Metropolitan	77 mins (n=2380)	514 mins (n=352)	82 mins (n=2732)
Rural	137 mins (n=618)	539 mins (n=297)	178 mins (n=915)
NSW overall	83 mins (n=3180)	546 mins (n=711)	96 mins (n=3891)

A number of major trauma patients (n=248, 6.0%) who arrived at a designated trauma service needed to be transferred to a higher level of care for specialised treatment. While vital, this transfer prolongs the time before the required (definitive) care can be provided, such as specialised surgery, interventional radiology or paediatric services. Overall, 904 major trauma patients were transferred from another acute care facility with a median time of injury to definitive care of 579 minutes (Table 20).*

Table 20: Median time of injury to arrival at definitive care (n=3950)

Location of injury	Direct from scene	Transferred from another hospital	Overall
Metropolitan	77 mins (n=2323)	520 mins (n=429)	84 mins (n=2752)
Rural	142 mins (n=544)	592 mins (n=390)	211 mins (n=934)
NSW overall	83 mins (n=3046)	579 mins (n=904)	103 mins (n=3950)

* See [Glossary](#) for definition of definitive care

Mode of transport

Major trauma patients are transported to a designated trauma service from either the scene of the injury or from another hospital. Road ambulance was by far the most common mode of transport (76.2%), with helicopter, fixed wing aircraft, and private transport also used (Table 21, Figures 20-21).

Table 21: Mode of transport to definitive care (n=4121)

Transport mode	Direct from scene of injury	Transfer from another acute care facility	Total
Road ambulance	2518 (79.2%)	624 (66.3%)	3142 (76.2%)
Helicopter	398 (12.5%)	150 (15.9%)	548 (13.3%)
Private vehicle	240 (7.5%)	6 (0.6%)	246 (6.0%)
Unknown	17 (0.5%)	23 (2.4%)	40 (1.0%)
Other	6 (0.2%)	19 (2.0%)	25 (0.6%)
Fixed wing aircraft	1 (0%)	119 (12.6%)	120 (2.9%)
Total	3180	941	4121

Figure 20: Mode of transport to definitive care when transported direct from the scene of injury, by injury location (n=2998)





Figure 21: Mode of transport to definitive care when transferred from another acute care facility, by injury location (n=854)



Major trauma admissions


Introduction

The information in this section of the report is based on the number of major trauma patients admitted to NSW trauma services (n=4252), not the number of major trauma patients (n=4140), as some patients were treated in more than one NSW reporting facility. Trauma patient data is discussed in detail in the [Major trauma patients section](#).

Overall burden of trauma

Major trauma patients are an important cohort, but only represent a fraction of all trauma patients who are admitted to facilities across NSW. Based on data from the Admitted Patient Data Collection (APDC) and using an International Classification of Diseases (ICD) to AIS mapping tool, 30,147 patients with an estimated ISS >5 were admitted to a NSW health facility during the reporting period (Figure 22).^{13,14} This demonstrates the significant burden that trauma, as a whole, places on the health system.

Figure 22: Number of patients with an estimated ISS >5 by local health district (n=30,147)



Facility overview

Table 22 provides an overview of the major trauma admissions for each designated trauma service together with averages from the three trauma service types.

Table 22: Overview of trauma service admissions (n=4252)

Facility	Number of admissions	Number of admissions (ISS >12)	Average age	Average ISS	Case fatality rate (ISS >12) excluding traumatic DOA
Adult major trauma services	3172	2683	55.0	18	9.2%
John Hunter Hospital	596	558	53.4	20	8.9%
Liverpool Hospital	585	432	55.5	16	8.8%
Royal North Shore Hospital	583	494	57.5	20	11.4%
Royal Prince Alfred Hospital	319	272	54.9	18	8.9%
St George Hospital	292	227	52.9	19	6.6%
St Vincent's Hospital	218	194	53.3	19	8.2%
Westmead Hospital	579	506	55.6	17	9.5%
Paediatric major trauma services	217	178	7.4	19	5.7%
John Hunter Children's Hospital	48	43	8.4	19	4.7%
Sydney Children's Hospital	72	51	6.8	15	2.0%
The Children's Hospital at Westmead	97	84	7.4	21	8.5%
Regional trauma services	863	715	54.4	17	6.3%
Coffs Harbour Base Hospital	76	63	50.0	20	6.3%
Gosford Hospital	54	46	58.3	15	8.7%
Lismore Base Hospital	78	73	56.9	19	13.7%
Nepean Hospital	120	67	60.2	13	9.0%
Orange Base Hospital	111	90	49.9	16	3.3%
Port Macquarie Base Hospital	67	57	47.3	18	1.8%
Tamworth Base Hospital	91	72	49.9	16	2.8%
Tweed Heads District Hospital	46	39	55.7	18	5.3%
Wagga Wagga Base Hospital	81	74	51.1	16	2.7%
Wollongong Hospital	139	134	60.3	18	8.3%

Admission type

Most major trauma patients (78.5%, n=3102) were admitted to a trauma service direct from the scene of injury. However, there were differences in the admission type between adult and paediatric patient groups (Table 23).


Table 23: Number of admissions by type (n=4252)

Admission type	Number of admissions (paediatric / adult)	Percentage of admissions (paediatric / adult)
Direct from scene	189 / 3102	63.4% / 78.5%
Transfer from acute care facility	108 / 834	36.2% / 21.1%
Unknown and others	1 / 18	0.3% / 0.5%

Revised Trauma Score

The Revised Trauma Score* is an early (<24 hours) indicator of trauma outcomes. The lower the score, the higher is the likelihood of death. There was a negative correlation between the Revised Trauma Score and ISS, showing that the higher the ISS, the lower the Revised Trauma Score. The average Revised Trauma Score was 7.08 (Figure 23).

Figure 23: Average Revised Trauma Score by ISS group (n=4000)




* See [Glossary](#) for a definition of the Revised Trauma Score.⁹

Trauma team activation


Major trauma patients require rapid, thorough and systematic assessment and resuscitation. This is conducted on arrival by a multidisciplinary team known as a trauma team. The activation of the trauma team is based on a locally derived set of criteria including the mechanism of injury and physiological parameters. 49.9% of patients who arrived at a trauma service direct from scene received a full trauma team activation compared to 19.6% of those who were transferred from another acute care facility (Figure 24).

Figure 24: Trauma team activation by admission type (n=4252)



As expected, the activation of the trauma team increased with severity of injury with 37.2% (n=366) of the ISS 13-15 group receiving a full trauma team activation compared to 79.2% (n=95) of the ISS 41-75 group (Figure 25).

Figure 25: Trauma team activation by ISS group (n=4243)



Vital signs on arrival to the emergency department

The initial vital signs on arrival to the emergency department (ED) are useful indicators of injury severity and predictors of death. Major trauma patients who presented with hypotension (systolic blood pressure <100mmHg), a Glasgow Coma Scale <13 or hypothermia (temperature <35°C) represented a higher proportion of those who died compared to all patients (Figures 26-28).

Figure 26: Initial systolic blood pressure on arrival to the ED and mortality (n= 4102)




Figure 27: Initial Glasgow Coma Scale on arrival to the ED and mortality (n=4,069)





Figure 28: Initial temperature on arrival to ED and mortality (n=3,646)




Intensive care unit admissions

Overall, 47.3% (n=2006) of major trauma patients received an intensive care unit (ICU) admission, with the percentage increasing with the level of injury severity (Figure 29). This is up from 42.2% the previous year. While the number of ICU admissions may be explained by the level of severity of injury, the complexity of injury and other factors such as comorbidities may also influence the requirement for ICU admission.

It is important to note that the ISS <13 group is only included in the report if they have been admitted to ICU or died, hence the high proportion of ICU admissions in this group.

Figure 29: ICU admission by ISS group (n=4243)



Length of stay

The length of stay (LOS) in hospital, particularly the LOS in an ICU, is an indicator of the severity and complexity of the injury. The average hospital LOS for major trauma patients was 12.8 days (Table 24).

Table 24: Overview of ICU and hospital length of stay (n=4252)

LOS description	Number of patients	Average days	Median days
ICU LOS	2008	5.0	2.0
Total hospital LOS	4252	12.8	6.0

The average ICU LOS increased with severity, with ISS 13-15 group being 4.5 days and ISS 41-75 group being 13.3 days. A similar increasing pattern was also shown in the hospital length of stay as expected (Table 25).

Table 25: ICU and hospital length of stay by ISS (n=4252)

ISS group	Average ICU LOS	Median ICU LOS	Average hospital LOS	Median hospital LOS
ISS <13	2.6	1.0	12.3	6.0
ISS 13-15	4.5	2.0	9.1	5.0
ISS 16-24	4.4	2.0	11.8	7.0
ISS 25-40	7.3	4.0	16.9	9.0
ISS 41-75	13.3	7.0	29.8	13.5

No considerable change was observed from the previous year in the median values of ICU LOS and hospital LOS across different age groups (Table 26).

Table 26: ICU and hospital length of stay by age (n=4252)

Age group	Average ICU LOS	Median ICU LOS	Average hospital LOS	Median hospital LOS
0-4	11.4	2	9.6	2
5-9	6.1	2	10.2	5
10-14	2.6	1	11.1	5
15-19	4.8	2	11.4	5
20-24	4.7	2	11.9	6
25-29	4.2	2	10.2	5
30-34	3.9	2	10.7	5
35-39	6	2	13.7	6
40-44	6.3	2	14.9	7
45-49	5.7	2	12.3	5.5
50-54	5.5	3	17.4	7
55-59	4.1	2	12.5	6
60-64	6.2	3	14.4	7
65-69	7.1	3	15.8	6
70-74	4.8	3	13.5	7
75-79	4.8	3	13.3	8.5
80-84	3.4	2	11.8	8
85+	3.1	2	11.7	7

Procedures


Despite the serious nature of the injuries sustained by major trauma patients, the number of surgical or interventional radiology procedures performed was very low. There were 841 procedures performed on 754 patients (17.7% of all admissions). The highest percentage of procedures were performed in the ISS 41-75 group (Table 27, Figure 30).

Table 27: Procedures performed by type (n=754)

Procedure	Number of procedures	Percentage of total admissions
Open reduction internal fixation	399	9.4%
Laparotomy	176	4.1%
Craniotomy	129	3.0%
Embolisation	99	2.3%
Thoracotomy	38	0.9%
Total	841	17.7%


An increase in the proportion of laparotomy, open reduction internal fixation and embolisation procedures performed is noted in the ISS 41-75 group (Figure 30).

Figure 30: Procedures performed by ISS group (n=839)



Over the last five years, there has not been a substantial change in the number of procedures performed, with only embolisation procedures showing a notable increase (Figure 31).


Figure 31: Five-year trend of procedures performed (n=3985)



Discharge destination of survivors


Survivors of major trauma (n= 3846) were discharged to various locations, with the home being the most common (65.3%) followed by rehabilitation (15.1%). 'Other' incorporates locations such as board and care, burns centre, foster care, residential institution, missing and unknown (Figure 32).

Figure 32: Discharge destination of survivors (n=3846)



Overall, the rate of discharge home decreased as the injury severity increased, coinciding with an increase in the rate of discharge to rehabilitation services (Figure 33).

Figure 33: Discharge destination of survivors by ISS group (n=3844)



References

1. NSW Trauma Registry. NSW Agency for Clinical Innovation; 2020.
2. NSW Department of Health. [NSW Trauma Services Plan](#). Sydney: NSW Dept of Health; 2009.
3. NSW Agency for Clinician Innovation. [Improving the Quality of Trauma Care in NSW: Trauma Services Model of Care](#). Sydney: ACI; 2019.
4. Statistical Information Management Committee. Guidelines for the Use and Disclosure of Health Data for Statistical Purposes. Canberra: Australian Institute of Health and Welfare; 2007.
5. Centre for Epidemiology and Evidence. [Privacy issues and the reporting of small numbers](#). Sydney: NSW Ministry of Health; 2015.
6. Australian Bureau of Statistics. [Australian statistical geography standard \(ASGS\) remoteness structure](#). Canberra: Commonwealth of Australia; 2016.
7. Australia Government Department of Health. [Accessibility and Remoteness Index of Australia](#). Canberra: Dept of Health; 2006.
8. Association of the Advancement of Automotive Medicine. Abbreviated Injury Scale 2005 (Update 2008). Gennarelli T, Woodzin E (eds). Chicago, IL: 2008.
9. Champion HR, Sacco WJ, Copes WS, et al. A Revision of the Trauma Score. J Trauma. 1989 May;29:623-629. DOI: [10.1097/00005373-198905000-00017](#)
10. Australian Bureau of Statistics. [Standard Population for Use in Age-Standardisation Table](#). Australian Demographic Statistics. Catalogue No. 3101.0. Canberra: ABS; 2013.
11. Australian Bureau of Statistics. [Australian Demographic Statistics, Dec 2019](#). Catalogue No. 3101.0. Canberra: ABS; 2019.
12. Butcher N, Balogh Z. AIS >2 in at least two body regions: A potential new anatomical definition of polytrauma. Injury. 2012;43(2):196-199.
13. Admitted Patient Data Collection. NSW Ministry of Health; 2020.
14. Dinh MM, Singh H, Sarrami P, Levesque JF. Correlating injury severity scores and major trauma volume using a state-wide in-patient administrative dataset linked to trauma registry data — A retrospective analysis from New South Wales Australia. Injury. 2019 Sep 13; pii S0020-1383(19)30539-X DOI: [10.1016/j.injury.2019.09.022](#)

Glossary

Abbreviated Injury Scale (AIS) is an anatomically based, consensus-derived, global severity scoring system that classifies each injury by body region, according to its relative importance on a six-point ordinal scale. The AIS is the basis for the Injury Severity Score (ISS) calculation of the multiply injured patient.

Age-specific rate is an annualised rate given as per 100,000 persons, based on the estimated NSW population during the reporting period (end of December). Source: Australian Bureau of Statistics. Australian Demographic Statistics, Catalogue No. 3101.0. Canberra: ABS.¹¹

Age-standardised rate is an annualised rate given as per 100,000 persons, standardised to the Australian population at 30 June 2001. Source: Australian Bureau of Statistics. Standard Population for Use in Age-Standardisation Table. Australian Demographic Statistics, Mar 2013. Catalogue No. 3101. Canberra: ABS, Dec 2013.¹⁰

Case fatality rate is the proportion of deaths for a designated population expressed as a percentage. The NSW Trauma Minimum Data Set does not include all survivors with an ISS of less than or equal to 12. The case fatality rate in this report is only calculated on patients with an ISS of greater than 12 and will be expressed as 'case fatality rate for ISS >12'.

Definitive care is defined as the hospital providing the highest level of care to meet all the clinical needs of the patient. Many patients receive definitive care at regional trauma services, but a small number of patients are transferred to a major trauma service (higher level) for specialised care.

Geriatric population is defined as those aged 65 years or older.

Location of injury is defined as either metropolitan or rural based on the recorded postcode of injury. The process used to define the two categories is outlined in the [Methodology](#) section.

Injury Severity Score (ISS) assesses the combined effects of the multiply injured patient and is based on an anatomical injury severity classification, the AIS. The ISS is an internationally recognised scoring system which correlates with mortality, morbidity and other measures of severity. The ISS is calculated as the sum of the squares of the highest AIS code in each of the three most severely injured ISS body regions.

ISS body regions consist of six anatomical regions as defined in the AIS dictionary:

- head or neck
- face
- chest
- abdominal or pelvic contents
- extremities or pelvic girdle
- external.

Isolated fractured neck of femur is defined as the AIS codes 853161.3 and 853162.3 and where no other injury is recorded.

Major trauma is defined as all patients of any age, who were admitted to a designated NSW trauma service within seven days of sustaining an injury, and:

- had an ISS >12 (moderate to critically injured), or
- were admitted to an intensive care unit (irrespective of ISS) following injury, or

- died in hospital (irrespective of ISS) following injury, except those with an isolated fractured neck of femur injury sustained from a fall from a standing height (<1 metre) and those aged 65 years or older who die with minor soft tissue injury only.

Major trauma services can provide the full spectrum of care for major and moderately injured patients, from initial resuscitation through to rehabilitation and discharge. There are currently seven adult and three paediatric designated major trauma services in NSW.

Mechanism of injury refers to the mechanisms whereby energy is transferred from the environment to the person.

Minor soft tissue injury is defined as a superficial injury including abrasions, contusions, and lacerations (AIS codes: 910000.1; 910200.1; 910400.1; 910600.1, 810099.1, 810202.1, 810402.1, 810600.1, 810602.1, 710099.1, 710202.1, 710402.1, 710600.1, 710602.1, 510099.1, 510202.1, 510402.1, 510600.1, 510602.1, 410099.1, 410202.1, 410402.1, 410600.1, 410602.1, 310099.1, 310202.1, 310402.1, 310600.1, 310602.1, 210099.1, 210202.1, 210402.1, 210600.1, 210602.1).

Not further specified is an injury descriptor used in AIS coding where detailed information is lacking including injury type or severity.

Other transport incident is defined as a patient involved in an accident involving a device designed primarily for, or being used at the time primarily for, conveying persons or goods from one place to another (ICD 10 codes: V00 – V99 inclusive) that did not meet place of occurrence road trauma criteria.

Polytrauma is defined as serious injury (AIS severity >2) in two or more ISS body regions.

Regional trauma services can provide all aspects of care to patients with minor to moderate trauma, and definitive care to a limited number of major trauma patients in collaboration with the major trauma service. A regional trauma service provides initial assessment, stabilisation, definitive care and initiates transfer to a major trauma service when a patient requires services not available at the regional trauma service. There are currently ten designated regional trauma services in NSW.

Road trauma is defined as a patient involved in an accident involving a device designed primarily for, or being used at the time primarily for, conveying persons or goods from one place to another (ICD 10 codes: V00 – V89 inclusive) AND had a street, highway and other paved roadways as the place of occurrence of the external cause (ICD 10 codes: Y92.4 inclusive).

Revised Trauma Score is a physiological scoring system used for predicting death. It consists of the first set of vital signs data obtained on the patient after arrival at hospital including Glasgow Coma Scale, systolic blood pressure and respiratory rate. Values for the Revised Trauma Score are in the range 0 to 7.8408. The lower the score, the higher the likelihood of death.⁹

Standardised mortality ratio is a ratio between the observed number of deaths in a study population and the number of deaths that would be expected, based on the age or ISS specific rates in a standard population and the age or ISS distribution of the study population.

Traumatic death on arrival is defined as a patient presenting to the emergency department with no recordable pulse or blood pressure (pulse rate = 0; systolic blood pressure = 0), no motor response elicited during the Glasgow Coma Scale (GCS) assessment (GCS motor score = 1), and the post emergency disposition recorded as 'died'.

Abbreviations

AIS	Abbreviated Injury Scale
ASGS-RA	Australian Statistical Geography Standard Remoteness Areas
CCLHD	Central Coast Local Health District
DBP	Diastolic blood pressure
DOA	Dead on arrival
ED	Emergency department
FWLHD	Far West Local Health District
HNELHD	Hunter New England Local Health District
ICD	International Classification of Diseases
ICU	Intensive care unit
ISLHD	Illawarra Shoalhaven Local Health District
ISS	Injury Severity Score
ITIM	NSW Institute of Trauma and Injury Management
JH&FMHN	Justice Health & Forensic Mental Health Service Network
LOS	Length of stay
MLHD	Murrumbidgee Local Health District
MNCLHD	Mid North Coast Local Health District
MTS	Major trauma service
NBMLHD	Nepean Blue Mountains Local Health District
NNSWLHD	Northern New South Wales Local Health District
NSLHD	Northern Sydney Local Health District
PH	Prehospital
PTS	Paediatric trauma service
RTS	Regional trauma service or Revised Trauma Score
SaO ₂	Arterial oxygen saturation
SBP	Systolic blood pressure
SCHN	Sydney Children's Hospital Network
SESLHD	South Eastern Sydney Local Health District
SLHD	Sydney Local Health District
SNSWLHD	Southern New South Wales Local Health District
SVHN	St. Vincent's Health Network Local Health District
SWSLHD	South Western Sydney Local Health District
TRISS	Trauma and injury severity score
WNSWLHD	Western New South Wales Local Health District
WSLHD	Western Sydney Local Health District

Acknowledgements

The NSW Institute of Trauma and Injury Management (ITIM) wishes to acknowledge the NSW Trauma Services for their contribution of data to the NSW Trauma Registry.

ACI team

- Hardeep Singh, Data Officer, NSW ITIM
- Emma Curtin, TORQUE Support Nurse, NSW ITIM
- Glenn Sisson, Manager, NSW ITM
- Michael Dinh, Clinical Director, NSW ITIM
- Kelly Dee, Clinical Review Officer, NSW ITIM
- Jon Newman, Project Officer, NSW ITIM
- Pooria Sarrami, Research Officer, NSW ITIM
- Chelsea McBride, Project Officer, NSW ITIM

Appendices


List of appendices

- [Appendix 1: Data completeness](#)
- [Appendix 2: Adult major trauma service summaries](#)
- [Appendix 3: Paediatric major trauma service summaries](#)
- [Appendix 4: Regional trauma service summaries](#)
- [Appendix 5: Calculation of the Injury Severity Score](#)
- [Appendix 6: Australian Statistical Geography Standard Remoteness Areas](#)

Appendix 1: Data completeness

High levels of data accuracy and entry completion, as entered by the trauma facilities, is crucial in enabling accurate and thorough data analysis as part of a broader trauma quality assurance program. The average overall data completion of mandatory elements was 95.0%, with completion rates ranging from 90.4% to 99.7% (Figure 34).

Figure 34: Overall data completeness by facility (n=4252)



A detailed breakdown of data completion by facility, showing each mandatory data element can be seen in Table 28.

Table 28: Detailed data completeness by facility

Legend colour guide (Percentage complete)																				
100%				80-99%				60-79%				Less than 60%								
Facility	Coffs Harbour Base Hospital	Gosford Hospital	John Hunter Children's Hospital	John Hunter Hospital	Lismore Base Hospital	Liverpool Hospital	Nepean Hospital	Orange Base Hospital	Port Macquarie Base Hospital	Royal North Shore Hospital	Royal Prince Alfred Hospital	St George Hospital	St Vincent's Hospital	Sydney Children's Hospital	Tamworth Base Hospital	The Children's Hospital at Westmead	Tweed Heads District Hospital	Wagga Wagga Base Hospital	Westmead Hospital	Wollongong Hospital
Trauma ID																				
Age																				
Gender																				
Post code home																				
Post code injury																				
Injury date																				
Injury time																				
Primary cause of injury																				
Injury type																				
Injury place																				
Injury activity																				
Fall height																				
Scene agency																				
Scene mode																				
PH agency Run No.																				
PH agency call received date																				
PH agency call received time																				
PH agency arrive at patient date																				
PH agency arrive at patient time																				
PH agency depart location date																				
PH agency depart location time																				
Admission date																				
Admission time																				
System access																				
ED arrival date																				
ED arrival time																				
ED depart date																				
ED depart time																				

Facility	Coffs Harbour Base Hospital	Gosford Hospital	John Hunter Children's Hospital	John Hunter Hospital	Lismore Base Hospital	Liverpool Hospital	Nepean Hospital	Orange Base Hospital	Port Macquarie Base Hospital	Royal North Shore Hospital	Royal Prince Alfred Hospital	St George Hospital	St Vincent's Hospital	Sydney Children's Hospital	Tamworth Base Hospital	The Children's Hospital at Westmead	Tweed Heads District Hospital	Wagga Wagga Base Hospital	Westmead Hospital	Wollongong Hospital
Trauma response																				
Post ED disposition																				
Intubated																				
Intubated method																				
Resp assisted																				
Resp assisted type																				
Paralytic agent																				
Sedation																				
ED vitals-Heart rate																				
ED vitals-Resp rate																				
ED vitals-DBP																				
ED vitals-SBP																				
ED vitals-Temp																				
ED vitals-SAO2																				
ED vitals-GCS eye																				
ED vitals-GCS verbal																				
ED vitals-GCS motor																				
ED vitals-GCS total																				
ED vitals-RTS total																				
Ref hospital 1 arrival date																				
Ref hospital 1 arrival time																				
Ref hospital 1 transfer rationale																				
Ref hospital 1 procedures																				
Ref hospital 2 arrival date																				
Ref hospital 2 arrival time																				
Ref hospital 2 transfer rationale																				
Ref hospital 2 procedures																				
Inter facility 1 transfer agency																				
Inter facility 1 mode																				
Inter facility 2 transfer agency																				
Inter facility 2 mode																				
Location tracking depts																				
Location tracking arrival dates																				
Location tracking arrival times																				
Procedure start dates																				
Procedure start times																				
AIS codes																				
ISS																				

Facility	Coffs Harbour Base Hospital	Gosford Hospital	John Hunter Children's Hospital	John Hunter Hospital	Lismore Base Hospital	Liverpool Hospital	Nepean Hospital	Orange Base Hospital	Port Macquarie Base Hospital	Royal North Shore Hospital	Royal Prince Alfred Hospital	St George Hospital	St Vincent's Hospital	Sydney Children's Hospital	Tamworth Base Hospital	The Children's Hospital at Westmead	Tweed Heads District Hospital	Wagga Wagga Base Hospital	Westmead Hospital	Wollongong Hospital
Outcome discharge disposition																				
Location of death																				
Discharge date																				
Discharge time																				
ICU length of stay																				
Hospital length of stay																				
Discharge destination																				
Discharge facility																				
TRISS																				
Record complete flag																				
Extrication minutes																				

Appendix 2: Adult major trauma service summaries

Table 29: Trauma data profile, John Hunter Hospital

Description	Facility	Peer
Total admissions	596	453.1
Mean monthly admissions	49.7	37.8
Case fatality rate (ISS >12 excl. traumatic DOA)	8.9%	9.2%
Sex		
Male / Female	435 / 161	323 / 131
Age ranges		
Mean age	53.4	55
0-4	0 (0%)	1.3 (0.3%)
5-9	0 (0%)	0.1 (0%)
10-14	0 (0%)	0.3 (0.1%)
15-19	33 (5.5%)	20.1 (4.4%)
20-24	43 (7.2%)	33.7 (7.4%)
25-29	30 (5%)	28 (6.2%)
30-34	33 (5.5%)	25 (5.5%)
35-39	44 (7.4%)	27.4 (6.1%)
40-44	35 (5.9%)	24.6 (5.4%)
45-49	42 (7%)	28.3 (6.2%)
50-54	47 (7.9%)	28.4 (6.3%)
55-59	40 (6.7%)	29.9 (6.6%)
60-64	49 (8.2%)	33.7 (7.4%)
65-69	37 (6.2%)	27.1 (6%)
70-74	40 (6.7%)	29.9 (6.6%)
75-79	45 (7.6%)	30.6 (6.7%)
80-84	31 (5.2%)	33.6 (7.4%)
85+	47 (7.9%)	51.1 (11.3%)
Injury Severity Score		
Mean ISS	20.8	18.8
ISS <13	36 (6.1%)	69 (15.3%)
ISS 13-15	106 (17.8%)	102.6 (22.7%)
ISS 16-24	270 (45.5%)	167.7 (37.1%)
ISS 25-40	152 (25.6%)	99.4 (22%)
ISS 41-75	30 (5.1%)	13.6 (3%)
Mechanism of injury		
Assault	42 (7%)	29.1 (6.4%)
Falls	225 (37.8%)	206.3 (45.5%)
Falls ≥65 years (% of all mechanisms of injury for ≥65 years)	139 (69.5%)	128.9 (74.8%)
Road trauma	207 (34.7%)	133.7 (29.5%)
Other transport incident	56 (9.4%)	31.9 (7%)
All other injuries	66 (11.1%)	52.1 (11.5%)

Description	Facility	Peer
Injury type		
Blunt	546 (91.6%)	419.9 (92.7%)
Penetrating	35 (5.9%)	25.3 (5.6%)
Unknown	15 (2.5%)	8 (1.8%)
Admission type		
Direct admission	391 (65.6%)	353.3 (78%)
Transfer in	203 (34.1%)	98.4 (21.7%)
Unknown	2 (0.3%)	1.4 (0.3%)
Arrival modes		
Ambulance	398 (66.8%)	348.3 (76.9%)
Helicopter	126 (21.1%)	50.9 (11.2%)
Other (private vehicle, fixed wing aircraft, unknown)	72 (12.1%)	54 (11.9%)
Revised Trauma Score		
Mean overall	6.9	7
ISS <13	6.1	6.6
ISS 13-15	7.5	7.6
ISS 16-24	7.3	7.3
ISS 25-40	6.5	6.4
ISS 41-75	4.3	4.3
Hospital length of stay		
Total bed days	7867	6276.4
Mean overall	13.2	13.9
ISS <13	8.8	11.2
ISS 13-15	12.8	10
ISS 16-24	12.5	12.5
ISS 25-40	13.6	19.6
ISS 41-75	25.5	31.9
ICU length of stay		
ICU total bed days (number of ICU admissions)	1441 (214)	1200.6 (227.1)
Mean overall	6.7	5.3
ISS <13	2.8	2.7
ISS 13-15	3.6	4.1
ISS 16-24	6.3	4.6
ISS 25-40	8.2	8
ISS 41-75	10.7	11.1
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	907 (152)	765.9 (134.6)
Mean overall	6	5.7
ISS <13	3.2	2.8
ISS 13-15	2.9	4
ISS 16-24	6.1	4.5
ISS 25-40	5.8	7.7
ISS 41-75	9.2	10.7

Table 30: Trauma data profile, Liverpool Hospital

Description	Facility	Peer
Total admissions	585	453.1
Mean monthly admissions	48.8	37.8
Case fatality rate (ISS >12 excl. traumatic DOA)	8.8%	9.2%
Sex		
Male / Female	396 / 189	323 / 131
Age ranges		
Mean age	55.5	55
0-4	4 (0.7%)	1.3 (0.3%)
5-9	1 (0.2%)	0.1 (0%)
10-14	1 (0.2%)	0.3 (0.1%)
15-19	28 (4.8%)	20.1 (4.4%)
20-24	40 (6.8%)	33.7 (7.4%)
25-29	31 (5.3%)	28 (6.2%)
30-34	32 (5.5%)	25 (5.5%)
35-39	35 (6%)	27.4 (6.1%)
40-44	33 (5.6%)	24.6 (5.4%)
45-49	38 (6.5%)	28.3 (6.2%)
50-54	34 (5.8%)	28.4 (6.3%)
55-59	38 (6.5%)	29.9 (6.6%)
60-64	40 (6.8%)	33.7 (7.4%)
65-69	32 (5.5%)	27.1 (6%)
70-74	32 (5.5%)	29.9 (6.6%)
75-79	39 (6.7%)	30.6 (6.7%)
80-84	58 (9.9%)	33.6 (7.4%)
85+	69 (11.8%)	51.1 (11.3%)
Injury Severity Score		
Mean ISS	16.7	18.8
ISS <13	149 (25.6%)	69 (15.3%)
ISS 13-15	125 (21.5%)	102.6 (22.7%)
ISS 16-24	207 (35.6%)	167.7 (37.1%)
ISS 25-40	88 (15.1%)	99.4 (22%)
ISS 41-75	12 (2.1%)	13.6 (3%)
Mechanism of injury		
Assault	42 (7.2%)	29.1 (6.4%)
Falls	262 (44.8%)	206.3 (45.5%)
Falls ≥65 years (% of all mechanisms of injury for ≥65 years)	171 (74.3%)	128.9 (74.8%)
Road trauma	190 (32.5%)	133.7 (29.5%)
Other transport incident	39 (6.7%)	31.9 (7%)
All other injuries	52 (8.9%)	52.1 (11.5%)

Description	Facility	Peer
Injury type		
Blunt	558 (95.4%)	419.9 (92.7%)
Penetrating	27 (4.6%)	25.3 (5.6%)
Unknown	0 (0%)	8 (1.8%)
Admission type		
Direct admission	464 (79.3%)	353.3 (78%)
Transfer in	121 (20.7%)	98.4 (21.7%)
Unknown	0 (0%)	1.4 (0.3%)
Arrival modes		
Ambulance	452 (77.3%)	348.3 (76.9%)
Helicopter	36 (6.2%)	50.9 (11.2%)
Other (private vehicle, fixed wing aircraft, unknown)	97 (16.6%)	54 (11.9%)
Revised Trauma Score		
Mean overall	7	7
ISS <13	6.7	6.6
ISS 13-15	7.6	7.6
ISS 16-24	7.4	7.3
ISS 25-40	6.2	6.4
ISS 41-75	5.1	4.3
Hospital length of stay		
Total bed days	7114	6276.4
Mean overall	12.2	13.9
ISS <13	10.2	11.2
ISS 13-15	9.8	10
ISS 16-24	12.7	12.5
ISS 25-40	15.8	19.6
ISS 41-75	28.1	31.9
ICU length of stay		
ICU total bed days (number of ICU admissions)	1317 (356)	1200.6 (227.1)
Mean overall	3.7	5.3
ISS <13	2.2	2.7
ISS 13-15	3.3	4.1
ISS 16-24	3.6	4.6
ISS 25-40	7.9	8
ISS 41-75	8	11.1
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	633 (135)	765.9 (134.6)
Mean overall	4.7	5.7
ISS <13	2.7	2.8
ISS 13-15	2.9	4
ISS 16-24	3.8	4.5
ISS 25-40	7.9	7.7
ISS 41-75	6.9	10.7

Table 31: Trauma data profile, Royal North Shore Hospital

Description	Facility	Peer
Total admissions	583	453.1
Mean monthly admissions	48.6	37.8
Case fatality rate (ISS >12 excl. traumatic DOA)	11.4%	9.2%
Sex		
Male / Female	392 / 191	323 / 131
Age ranges		
Mean age	57.5	55
0-4	2 (0.3%)	1.3 (0.3%)
5-9	0 (0%)	0.1 (0%)
10-14	1 (0.2%)	0.3 (0.1%)
15-19	25 (4.3%)	20.1 (4.4%)
20-24	34 (5.8%)	33.7 (7.4%)
25-29	35 (6%)	28 (6.2%)
30-34	23 (3.9%)	25 (5.5%)
35-39	32 (5.5%)	27.4 (6.1%)
40-44	25 (4.3%)	24.6 (5.4%)
45-49	39 (6.7%)	28.3 (6.2%)
50-54	40 (6.9%)	28.4 (6.3%)
55-59	38 (6.5%)	29.9 (6.6%)
60-64	37 (6.3%)	33.7 (7.4%)
65-69	35 (6%)	27.1 (6%)
70-74	41 (7%)	29.9 (6.6%)
75-79	43 (7.4%)	30.6 (6.7%)
80-84	58 (9.9%)	33.6 (7.4%)
85+	75 (12.9%)	51.1 (11.3%)
Injury Severity Score		
Mean ISS	20.2	18.8
ISS <13	89 (15.3%)	69 (15.3%)
ISS 13-15	91 (15.6%)	102.6 (22.7%)
ISS 16-24	213 (36.5%)	167.7 (37.1%)
ISS 25-40	171 (29.3%)	99.4 (22%)
ISS 41-75	19 (3.3%)	13.6 (3%)
Mechanism of injury		
Assault	18 (3.1%)	29.1 (6.4%)
Falls	308 (52.8%)	206.3 (45.5%)
Falls ≥65 years (% of all mechanisms of injury for ≥65 years)	193 (76.6%)	128.9 (74.8%)
Road trauma	125 (21.4%)	133.7 (29.5%)
Other transport incident	45 (7.7%)	31.9 (7%)
All other injuries	87 (14.9%)	52.1 (11.5%)

Description	Facility	Peer
Injury type		
Blunt	533 (91.4%)	419.9 (92.7%)
Penetrating	16 (2.7%)	25.3 (5.6%)
Unknown	34 (5.8%)	8 (1.8%)
Admission type		
Direct admission	429 (73.6%)	353.3 (78%)
Transfer in	153 (26.2%)	98.4 (21.7%)
Unknown	1 (0.2%)	1.4 (0.3%)
Arrival modes		
Ambulance	447 (76.7%)	348.3 (76.9%)
Helicopter	72 (12.3%)	50.9 (11.2%)
Other (private vehicle, fixed wing aircraft, unknown)	64 (11%)	54 (11.9%)
Revised Trauma Score		
Mean overall	7	7
ISS <13	7.2	6.6
ISS 13-15	7.5	7.6
ISS 16-24	7.3	7.3
ISS 25-40	6.5	6.4
ISS 41-75	4.3	4.3
Hospital length of stay		
Total bed days	10841	6276.4
Mean overall	18.6	13.9
ISS <13	10.9	11.2
ISS 13-15	12.1	10
ISS 16-24	15.4	12.5
ISS 25-40	28.7	19.6
ISS 41-75	30.9	31.9
ICU length of stay		
ICU total bed days (number of ICU admissions)	1924 (378)	1200.6 (227.1)
Mean overall	5.1	5.3
ISS <13	2.2	2.7
ISS 13-15	1.6	4.1
ISS 16-24	3.6	4.6
ISS 25-40	8.4	8
ISS 41-75	10.1	11.1
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	1273 (195)	765.9 (134.6)
Mean overall	6.5	5.7
ISS <13	2.7	2.8
ISS 13-15	1.9	4
ISS 16-24	3.4	4.5
ISS 25-40	10	7.7
ISS 41-75	7.6	10.7

Table 32: Trauma data profile, Royal Prince Alfred Hospital

Description	Facility	Peer
Total admissions	319	453.1
Mean monthly admissions	26.6	37.8
Case fatality rate (ISS >12 excl. traumatic DOA)	8.9%	9.2%
Sex		
Male / Female	225 / 94	323 / 131
Age ranges		
Mean age	54.9	55
0-4	3 (0.9%)	1.3 (0.3%)
5-9	0 (0%)	0.1 (0%)
10-14	0 (0%)	0.3 (0.1%)
15-19	6 (1.9%)	20.1 (4.4%)
20-24	26 (8.2%)	33.7 (7.4%)
25-29	35 (11%)	28 (6.2%)
30-34	21 (6.6%)	25 (5.5%)
35-39	15 (4.7%)	27.4 (6.1%)
40-44	16 (5%)	24.6 (5.4%)
45-49	16 (5%)	28.3 (6.2%)
50-54	10 (3.1%)	28.4 (6.3%)
55-59	22 (6.9%)	29.9 (6.6%)
60-64	23 (7.2%)	33.7 (7.4%)
65-69	19 (6%)	27.1 (6%)
70-74	23 (7.2%)	29.9 (6.6%)
75-79	20 (6.3%)	30.6 (6.7%)
80-84	24 (7.5%)	33.6 (7.4%)
85+	40 (12.5%)	51.1 (11.3%)
Injury Severity Score		
Mean ISS	18.4	18.8
ISS < 13	47 (14.7%)	69 (15.3%)
ISS 13-15	69 (21.6%)	102.6 (22.7%)
ISS 16-24	127 (39.8%)	167.7 (37.1%)
ISS 25-40	68 (21.3%)	99.4 (22%)
ISS 41-75	8 (2.5%)	13.6 (3%)
Mechanism of injury		
Assault	31 (9.7%)	29.1 (6.4%)
Falls	169 (53%)	206.3 (45.5%)
Falls ≥65 years (% of all mechanisms of injury for ≥65 years)	106 (84.1%)	128.9 (74.8%)
Road trauma	68 (21.3%)	133.7 (29.5%)
Other transport incident	3 (0.9%)	31.9 (7%)
All other injuries	48 (15%)	52.1 (11.5%)

Description	Facility	Peer
Injury type		
Blunt	279 (87.5%)	419.9 (92.7%)
Penetrating	39 (12.2%)	25.3 (5.6%)
Unknown	1 (0.3%)	8 (1.8%)
Admission type		
Direct admission	276 (86.5%)	353.3 (78%)
Transfer in	43 (13.5%)	98.4 (21.7%)
Unknown	0 (0%)	1.4 (0.3%)
Arrival modes		
Ambulance	274 (85.9%)	348.3 (76.9%)
Helicopter	0 (0%)	50.9 (11.2%)
Other (private vehicle, fixed wing aircraft, unknown)	45 (14.1%)	54 (11.9%)
Revised Trauma Score		
Mean overall	7.1	7
ISS <13	6.4	6.6
ISS 13-15	7.7	7.6
ISS 16-24	7.5	7.3
ISS 25-40	6.7	6.4
ISS 41-75	4	4.3
Hospital length of stay		
Total bed days	2962	6276.4
Mean overall	9.3	13.9
ISS <13	7.8	11.2
ISS 13-15	5.3	10
ISS 16-24	7.8	12.5
ISS 25-40	14.4	19.6
ISS 41-75	33.1	31.9
ICU length of stay		
ICU total bed days (number of ICU admissions)	576 (149)	1200.6 (227.1)
Mean overall	3.9	5.3
ISS <13	2	2.7
ISS 13-15	2.9	4.1
ISS 16-24	3.6	4.6
ISS 25-40	5.3	8
ISS 41-75	9.2	11.1
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	276 (86)	765.9 (134.6)
Mean overall	3.2	5.7
ISS <13	1.8	2.8
ISS 13-15	2	4
ISS 16-24	3.1	4.5
ISS 25-40	4	7.7
ISS 41-75	6.9	10.7

Table 33: Trauma data profile, St George Hospital

Description	Facility	Peer
Total admissions	292	453.1
Mean monthly admissions	24.3	37.8
Case fatality rate (ISS >12 excl. traumatic DOA)	6.6%	9.2%
Sex		
Male / Female	219 / 73	323 / 131
Age ranges		
Mean age	52.9	55
0-4	0 (0%)	1.3 (0.3%)
5-9	0 (0%)	0.1 (0%)
10-14	0 (0%)	0.3 (0.1%)
15-19	21 (7.2%)	20.1 (4.4%)
20-24	22 (7.5%)	33.7 (7.4%)
25-29	20 (6.8%)	28 (6.2%)
30-34	18 (6.2%)	25 (5.5%)
35-39	14 (4.8%)	27.4 (6.1%)
40-44	16 (5.5%)	24.6 (5.4%)
45-49	20 (6.8%)	28.3 (6.2%)
50-54	20 (6.8%)	28.4 (6.3%)
55-59	12 (4.1%)	29.9 (6.6%)
60-64	23 (7.9%)	33.7 (7.4%)
65-69	20 (6.8%)	27.1 (6%)
70-74	23 (7.9%)	29.9 (6.6%)
75-79	25 (8.6%)	30.6 (6.7%)
80-84	12 (4.1%)	33.6 (7.4%)
85+	26 (8.9%)	51.1 (11.3%)
Injury Severity Score		
Mean ISS	19	18.8
ISS <13	65 (22.3%)	69 (15.3%)
ISS 13-15	63 (21.6%)	102.6 (22.7%)
ISS 16-24	88 (30.1%)	167.7 (37.1%)
ISS 25-40	61 (20.9%)	99.4 (22%)
ISS 41-75	15 (5.1%)	13.6 (3%)
Mechanism of injury		
Assault	12 (4.1%)	29.1 (6.4%)
Falls	115 (39.4%)	206.3 (45.5%)
Falls ≥65 years (% of all mechanisms of injury for ≥65 years)	71 (67%)	128.9 (74.8%)
Road trauma	111 (38%)	133.7 (29.5%)
Other transport incident	27 (9.2%)	31.9 (7%)
All other injuries	27 (9.2%)	52.1 (11.5%)

Description	Facility	Peer
Injury type		
Blunt	278 (95.2%)	419.9 (92.7%)
Penetrating	12 (4.1%)	25.3 (5.6%)
Unknown	2 (0.7%)	8 (1.8%)
Admission type		
Direct admission	226 (77.4%)	353.3 (78%)
Transfer in	66 (22.6%)	98.4 (21.7%)
Unknown	0 (0%)	1.4 (0.3%)
Arrival modes		
Ambulance	222 (76%)	348.3 (76.9%)
Helicopter	47 (16.1%)	50.9 (11.2%)
Other (private vehicle, fixed wing aircraft, unknown)	23 (7.9%)	54 (11.9%)
Revised Trauma Score		
Mean overall	6.8	7
ISS <13	6.6	6.6
ISS 13-15	7.7	7.6
ISS 16-24	7.2	7.3
ISS 25-40	6.3	6.4
ISS 41-75	4	4.3
Hospital length of stay		
Total bed days	5821	6276.4
Mean overall	19.9	13.9
ISS <13	16	11.2
ISS 13-15	10.6	10
ISS 16-24	15.5	12.5
ISS 25-40	29.4	19.6
ISS 41-75	63.5	31.9
ICU length of stay		
ICU total bed days (number of ICU admissions)	1301 (203)	1200.6 (227.1)
Mean overall	6.4	5.3
ISS <13	3.2	2.7
ISS 13-15	3.4	4.1
ISS 16-24	4.8	4.6
ISS 25-40	9.1	8
ISS 41-75	18.2	11.1
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	981 (128)	765.9 (134.6)
Mean overall	7.7	5.7
ISS <13	2.6	2.8
ISS 13-15	3.3	4
ISS 16-24	3.5	4.5
ISS 25-40	9.2	7.7
ISS 41-75	24.9	10.7

Table 34: Trauma data profile, St Vincent's Hospital

Description	Facility	Peer
Total admissions	218	453.1
Mean monthly admissions	18.2	37.8
Case fatality rate (ISS >12 excl. traumatic DOA)	8.2%	9.2%
Sex		
Male / Female	168 / 50	323 / 131
Age ranges		
Mean age	53.3	55
0-4	0 (0%)	1.3 (0.3%)
5-9	0 (0%)	0.1 (0%)
10-14	0 (0%)	0.3 (0.1%)
15-19	6 (2.8%)	20.1 (4.4%)
20-24	25 (11.5%)	33.7 (7.4%)
25-29	10 (4.6%)	28 (6.2%)
30-34	19 (8.7%)	25 (5.5%)
35-39	14 (6.4%)	27.4 (6.1%)
40-44	8 (3.7%)	24.6 (5.4%)
45-49	14 (6.4%)	28.3 (6.2%)
50-54	17 (7.8%)	28.4 (6.3%)
55-59	17 (7.8%)	29.9 (6.6%)
60-64	18 (8.3%)	33.7 (7.4%)
65-69	9 (4.1%)	27.1 (6%)
70-74	13 (6%)	29.9 (6.6%)
75-79	6 (2.8%)	30.6 (6.7%)
80-84	16 (7.3%)	33.6 (7.4%)
85+	26 (11.9%)	51.1 (11.3%)
Injury Severity Score		
Mean ISS	19	18.8
ISS <13	24 (11%)	69 (15.3%)
ISS 13-15	64 (29.4%)	102.6 (22.7%)
ISS 16-24	73 (33.5%)	167.7 (37.1%)
ISS 25-40	51 (23.4%)	99.4 (22%)
ISS 41-75	6 (2.8%)	13.6 (3%)
Mechanism of injury		
Assault	17 (7.8%)	29.1 (6.4%)
Falls	122 (56%)	206.3 (45.5%)
Falls ≥65 years (% of all mechanisms of injury for ≥65 years)	59 (84.3%)	128.9 (74.8%)
Road trauma	56 (25.7%)	133.7 (29.5%)
Other transport incident	2 (0.9%)	31.9 (7%)
All other injuries	21 (9.6%)	52.1 (11.5%)

Description	Facility	Peer
Injury type		
Blunt	205 (94%)	419.9 (92.7%)
Penetrating	9 (4.1%)	25.3 (5.6%)
Unknown	4 (1.8%)	8 (1.8%)
Admission type		
Direct admission	216 (99.1%)	353.3 (78%)
Transfer in	2 (0.9%)	98.4 (21.7%)
Unknown	0 (0%)	1.4 (0.3%)
Arrival modes		
Ambulance	199 (91.3%)	348.3 (76.9%)
Helicopter	0 (0%)	50.9 (11.2%)
Other (private vehicle, fixed wing aircraft, unknown)	19 (8.7%)	54 (11.9%)
Revised Trauma Score		
Mean overall	7.2	7
ISS <13	6.6	6.6
ISS 13-15	7.7	7.6
ISS 16-24	7.6	7.3
ISS 25-40	6.6	6.4
ISS 41-75	4	4.3
Hospital length of stay		
Total bed days	2262	6276.4
Mean overall	10.4	13.9
ISS <13	7.5	11.2
ISS 13-15	6.4	10
ISS 16-24	11.2	12.5
ISS 25-40	14.7	19.6
ISS 41-75	16.8	31.9
ICU length of stay		
ICU total bed days (number of ICU admissions)	415 (93)	1200.6 (227.1)
Mean overall	4.5	5.3
ISS <13	2	2.7
ISS 13-15	2.3	4.1
ISS 16-24	5.4	4.6
ISS 25-40	5.7	8
ISS 41-75	6.5	11.1
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	292 (64)	765.9 (134.6)
Mean overall	4.6	5.7
ISS <13	2.2	2.8
ISS 13-15	5	4
ISS 16-24	6.1	4.5
ISS 25-40	5.3	7.7
ISS 41-75	4.8	10.7

Table 35: Trauma data profile, Westmead Hospital

Description	Facility	Peer
Total admissions	579	453.1
Mean monthly admissions	48.2	37.8
Case fatality rate (ISS >12 excl. traumatic DOA)	9.5%	9.2%
Sex		
Male / Female	423 / 156	323 / 131
Age ranges		
Mean age	55.6	55
0-4	0 (0%)	1.3 (0.3%)
5-9	0 (0%)	0.1 (0%)
10-14	0 (0%)	0.3 (0.1%)
15-19	22 (3.8%)	20.1 (4.4%)
20-24	46 (7.9%)	33.7 (7.4%)
25-29	35 (6%)	28 (6.2%)
30-34	29 (5%)	25 (5.5%)
35-39	38 (6.6%)	27.4 (6.1%)
40-44	39 (6.7%)	24.6 (5.4%)
45-49	29 (5%)	28.3 (6.2%)
50-54	31 (5.4%)	28.4 (6.3%)
55-59	42 (7.3%)	29.9 (6.6%)
60-64	46 (7.9%)	33.7 (7.4%)
65-69	38 (6.6%)	27.1 (6%)
70-74	37 (6.4%)	29.9 (6.6%)
75-79	36 (6.2%)	30.6 (6.7%)
80-84	36 (6.2%)	33.6 (7.4%)
85+	75 (13%)	51.1 (11.3%)
Injury Severity Score		
Mean ISS	17.4	18.8
ISS <13	73 (12.6%)	69 (15.3%)
ISS 13-15	200 (34.5%)	102.6 (22.7%)
ISS 16-24	196 (33.9%)	167.7 (37.1%)
ISS 25-40	105 (18.1%)	99.4 (22%)
ISS 41-75	5 (0.9%)	13.6 (3%)
Mechanism of injury		
Assault	42 (7.3%)	29.1 (6.4%)
Falls	243 (42%)	206.3 (45.5%)
Falls ≥65 years (% of all mechanisms of injury for ≥65 years)	163 (73.4%)	128.9 (74.8%)
Road trauma	179 (30.9%)	133.7 (29.5%)
Other transport incident	51 (8.8%)	31.9 (7%)
All other injuries	64 (11.1%)	52.1 (11.5%)

Description	Facility	Peer
Injury type		
Blunt	540 (93.3%)	419.9 (92.7%)
Penetrating	39 (6.7%)	25.3 (5.6%)
Unknown	0 (0%)	8 (1.8%)
Admission type		
Direct admission	471 (81.3%)	353.3 (78%)
Transfer in	101 (17.4%)	98.4 (21.7%)
Unknown	7 (1.2%)	1.4 (0.3%)
Arrival modes		
Ambulance	446 (77%)	348.3 (76.9%)
Helicopter	75 (13%)	50.9 (11.2%)
Other (private vehicle, fixed wing, unknown)	58 (10%)	54 (11.9%)
Revised Trauma Score		
Mean - overall	6.9	7
ISS <13	6.1	6.6
ISS 13-15	7.4	7.6
ISS 16-24	7.1	7.3
ISS 25-40	6.4	6.4
ISS 41-75	4	4.3
Hospital length of stay		
Total bed days	7068	6276.4
Mean overall	12.2	13.9
ISS <13	13.8	11.2
ISS 13-15	10.2	10
ISS 16-24	11.6	12.5
ISS 25-40	16.5	19.6
ISS 41-75	5.4	31.9
ICU length of stay		
ICU total bed days (number of ICU admissions)	1430 (197)	1200.6 (227.1)
Mean overall	7.3	5.3
ISS <13	4.6	2.7
ISS 13-15	8.6	4.1
ISS 16-24	6.8	4.6
ISS 25-40	10.1	8
ISS 41-75	5	11.1
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	999 (182)	765.9 (134.6)
Mean overall	5.5	5.7
ISS <13	3.5	2.8
ISS 13-15	6.1	4
ISS 16-24	5.2	4.5
ISS 25-40	7.5	7.7
ISS 41-75	3.5	10.7

Appendix 3: Paediatric major trauma service summaries

Table 36: Trauma data profile, John Hunter Children's Hospital

Description	Facility	Peer
Total admissions	48	72.3
Mean monthly admissions	4	6
Case fatality rate (ISS >12 excl. traumatic DOA)	4.7%	4.5%
Sex		
Male / Female	37 / 11	48 / 24
Age ranges		
Mean age	8.4	7.4
0-4	16 (33.3%)	27.3 (37.8%)
5-9	9 (18.8%)	15 (20.7%)
10-14	15 (31.2%)	23 (31.8%)
15-19	8 (16.7%)	7 (9.7%)
Injury Severity Score		
Mean ISS	19.9	19.4
ISS <13	5 (10.4%)	13 (18%)
ISS 13-15	8 (16.7%)	13.3 (18.4%)
ISS 16-24	22 (45.8%)	25.3 (35%)
ISS 25-40	10 (20.8%)	16 (22.1%)
ISS 41-75	3 (6.2%)	4.7 (6.5%)
Mechanism of injury		
Assault	3 (6.2%)	3.7 (5.1%)
Falls	8 (16.7%)	21 (29%)
Road trauma	10 (20.8%)	13.7 (18.9%)
Other transport incident	13 (27.1%)	12.3 (17.1%)
All other injuries	14 (29.2%)	21.7 (30%)
Injury type		
Blunt	39 (81.2%)	60.7 (83.9%)
Penetrating	2 (4.2%)	3.7 (5.1%)
Unknown	7 (14.6%)	8 (11.1%)
Admission type		
Direct admission	30 (62.5%)	38.3 (53%)
Transfer in	18 (37.5%)	33.7 (46.5%)
Unknown	0 (0%)	0.3 (0.5%)
Arrival modes		
Ambulance	27 (56.2%)	38 (52.5%)
Helicopter	9 (18.8%)	9.7 (13.4%)
Other (private vehicle, fixed wing aircraft, unknown)	12 (25%)	24.7 (34.1%)

Description	Facility	Peer
Revised Trauma Score		
Mean overall	6.6	6.6
ISS <13	7	6.3
ISS 13-15	7.6	7.3
ISS 16-24	6.7	7
ISS 25-40	6.8	6.2
ISS 41-75	2.8	3.9
Hospital length of stay		
Total bed days	483	872.3
Mean overall	10.1	12.1
ISS <13	2.2	7.1
ISS 13-15	7.1	5.9
ISS 16-24	13.2	12.4
ISS 25-40	8.3	13.4
ISS 41-75	14	38
ICU length of stay		
ICU total bed days (number of ICU admissions)	114 (28)	282.7 (42)
Mean overall	4.1	6.7
ISS <13	0.8	1.8
ISS 13-15	32	7.4
ISS 16-24	3.6	5.1
ISS 25-40	3.2	6.2
ISS 41-75	4.5	28.4
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	46 (13)	228.3 (28.3)
Mean overall	3.5	8.1
ISS <13	2	3.5
ISS 13-15	0	3.5
ISS 16-24	3	4.7
ISS 25-40	6.5	6.6
ISS 41-75	3.3	27.7

Table 37: Trauma data profile, Sydney Children's Hospital

Description	Facility	Peer
Total admissions	72	72.3
Mean monthly admissions	6	6
Case fatality rate (ISS >12 excl. traumatic DOA)	2%	4.5%
Sex		
Male / Female	51 / 21	48 / 24
Age ranges		
Mean age	6.8	7.4
0-4	31 (43.1%)	27.3 (37.8%)
5-9	13 (18.1%)	15 (20.7%)
10-14	24 (33.3%)	23 (31.8%)
15-19	4 (5.6%)	7 (9.7%)
Injury Severity Score		
Mean ISS	15.9	19.4
ISS <13	21 (29.2%)	13 (18%)
ISS 13-15	22 (30.6%)	13.3 (18.4%)
ISS 16-24	16 (22.2%)	25.3 (35%)
ISS 25-40	11 (15.3%)	16 (22.1%)
ISS 41-75	2 (2.8%)	4.7 (6.5%)
Mechanism of injury		
Assault	5 (6.9%)	3.7 (5.1%)
Falls	30 (41.7%)	21 (29%)
Road trauma	9 (12.5%)	13.7 (18.9%)
Other transport incident	11 (15.3%)	12.3 (17.1%)
All other injuries	17 (23.6%)	21.7 (30%)
Injury type		
Blunt	70 (97.2%)	60.7 (83.9%)
Penetrating	0 (0%)	3.7 (5.1%)
Unknown	2 (2.8%)	8 (11.1%)
Admission type		
Direct admission	26 (36.1%)	38.3 (53%)
Transfer in	46 (63.9%)	33.7 (46.5%)
Unknown	0 (0%)	0.3 (0.5%)
Arrival modes		
Ambulance	30 (41.7%)	38 (52.5%)
Helicopter	5 (6.9%)	9.7 (13.4%)
Other (private vehicle, fixed wing aircraft, unknown)	37 (51.4%)	24.7 (34.1%)

Description	Facility	Peer
Revised Trauma Score		
Mean overall	6.7	6.6
ISS <13	6.3	6.3
ISS 13-15	7.2	7.3
ISS 16-24	7.3	7
ISS 25-40	6.3	6.2
ISS 41-75	4.9	3.9
Hospital length of stay		
Total bed days	525	872.3
Mean overall	7.3	12.1
ISS <13	6.1	7.1
ISS 13-15	3.9	5.9
ISS 16-24	10.6	12.4
ISS 25-40	10.2	13.4
ISS 41-75	14.5	38
ICU length of stay		
ICU total bed days (number of ICU admissions)	90 (40)	282.7 (42)
Mean overall	2.2	6.7
ISS <13	1.2	1.8
ISS 13-15	1.7	7.4
ISS 16-24	2.2	5.1
ISS 25-40	4	6.2
ISS 41-75	6	28.4
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	51 (20)	228.3 (28.3)
Mean overall	2.5	8.1
ISS <13	1.3	3.5
ISS 13-15	2	3.5
ISS 16-24	8	4.7
ISS 25-40	2	6.6
ISS 41-75	9	27.7

Table 38: Trauma data profile, The Children's Hospital at Westmead

Description	Facility	Peer
Total admissions	97	72.3
Mean monthly admissions	8.1	6
Case fatality rate (ISS >12 excl. traumatic DOA)	6.1%	4.5%
Sex		
Male / Female	56 / 41	48 / 24
Age ranges		
Mean age	7.4	7.4
0-4	35 (36.1%)	27.3 (37.8%)
5-9	23 (23.7%)	15 (20.7%)
10-14	30 (30.9%)	23 (31.8%)
15-19	9 (9.3%)	7 (9.7%)
Injury Severity Score		
Mean ISS	21.7	19.4
ISS <13	13 (13.4%)	13 (18%)
ISS 13-15	10 (10.3%)	13.3 (18.4%)
ISS 16-24	38 (39.2%)	25.3 (35%)
ISS 25-40	27 (27.8%)	16 (22.1%)
ISS 41-75	9 (9.3%)	4.7 (6.5%)
Mechanism of injury		
Assault	3 (3.1%)	3.7 (5.1%)
Falls	25 (25.8%)	21 (29%)
Road trauma	22 (22.7%)	13.7 (18.9%)
Other transport incident	13 (13.4%)	12.3 (17.1%)
All other injuries	34 (35.1%)	21.7 (30%)
Injury type		
Blunt	73 (75.3%)	60.7 (83.9%)
Penetrating	9 (9.3%)	3.7 (5.1%)
Unknown	15 (15.5%)	8 (11.1%)
Admission type		
Direct admission	59 (60.8%)	38.3 (53%)
Transfer in	37 (38.1%)	33.7 (46.5%)
Unknown	1 (1%)	0.3 (0.5%)
Arrival modes		
Ambulance	57 (58.8%)	38 (52.5%)
Helicopter	15 (15.5%)	9.7 (13.4%)
Other (private vehicle, fixed wing aircraft, unknown)	25 (25.8%)	24.7 (34.1%)

Description	Facility	Peer
Revised Trauma Score		
Mean overall	6.5	6.6
ISS <13	6.1	6.3
ISS 13-15	7.3	7.3
ISS 16-24	7	7
ISS 25-40	6	6.2
ISS 41-75	4.2	3.9
Hospital length of stay		
Total bed days	1609	872.3
Mean overall	16.8	12.1
ISS <13	10.4	7.1
ISS 13-15	9.2	5.9
ISS 16-24	12.8	12.4
ISS 25-40	16.7	13.4
ISS 41-75	51.2	38
ICU length of stay		
ICU total bed days (number of ICU admissions)	644 (58)	282.7 (42)
Mean overall	11.1	6.7
ISS <13	3.2	1.8
ISS 13-15	5	7.4
ISS 16-24	7.1	5.1
ISS 25-40	8.6	6.2
ISS 41-75	40	28.4
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	588 (52)	228.3 (28.3)
Mean overall	11.3	8.1
ISS <13	5.9	3.5
ISS 13-15	4	3.5
ISS 16-24	5	4.7
ISS 25-40	8	6.6
ISS 41-75	39.1	27.7

Appendix 4: Regional trauma service summaries

Table 39: Trauma data profile, Coffs Harbour Base Hospital

Description	Facility	Peer
Total admissions	76	86.3
Mean monthly admissions	6.3	7.2
Case fatality rate (ISS >12 excl. traumatic DOA)	6.3%	6.3%
Sex		
Male / Female	59 / 17	62 / 25
Age ranges		
Mean age	50	54.4
0-4	0 (0%)	1.1 (1.3%)
5-9	0 (0%)	0.9 (1%)
10-14	1 (1.3%)	1.1 (1.3%)
15-19	5 (6.6%)	4 (4.6%)
20-24	6 (7.9%)	6 (7%)
25-29	8 (10.5%)	4.7 (5.4%)
30-34	3 (3.9%)	3.5 (4.1%)
35-39	3 (3.9%)	2.9 (3.4%)
40-44	3 (3.9%)	5.2 (6%)
45-49	6 (7.9%)	6.2 (7.2%)
50-54	9 (11.8%)	6.7 (7.8%)
55-59	4 (5.3%)	5.7 (6.6%)
60-64	8 (10.5%)	5.6 (6.5%)
65-69	6 (7.9%)	4.8 (5.6%)
70-74	2 (2.6%)	5.8 (6.7%)
75-79	3 (3.9%)	5 (5.8%)
80-84	2 (2.6%)	7.6 (8.8%)
85+	7 (9.2%)	9.5 (11%)
Injury Severity Score		
Mean ISS	20.3	17.3
ISS <13	13 (17.1%)	14.5 (16.9%)
ISS 13-15	11 (14.5%)	22.6 (26.3%)
ISS 16-24	30 (39.5%)	32.7 (38%)
ISS 25-40	18 (23.7%)	15.1 (17.6%)
ISS 41-75	4 (5.3%)	1.1 (1.3%)
Mechanism of injury		
Assault	4 (5.3%)	5.5 (6.4%)
Falls	20 (26.3%)	34.2 (39.6%)
Falls ≥65 years (% of all mechanisms of injury for ≥65 years)	9 (45%)	23.1 (70.6%)
Road trauma	35 (46.1%)	25.1 (29.1%)
Other transport incident	7 (9.2%)	12.1 (14%)
All other injuries	10 (13.2%)	9.4 (10.9%)

Description	Facility	Peer
Injury type		
Blunt	69 (90.8%)	81 (93.9%)
Penetrating	6 (7.9%)	4.1 (4.8%)
Unknown	1 (1.3%)	1.2 (1.4%)
Admission type		
Direct admission	73 (96.1%)	70.3 (81.5%)
Transfer in	3 (3.9%)	15.2 (17.6%)
Unknown	0 (0%)	0.8 (0.9%)
Arrival modes		
Ambulance	68 (89.5%)	68.5 (79.4%)
Helicopter	3 (3.9%)	4.5 (5.2%)
Other (private vehicle, fixed wing aircraft, unknown)	5 (6.6%)	13.3 (15.4%)
Revised Trauma Score		
Mean overall	7.4	7.5
ISS <13	7.7	7.6
ISS 13-15	7.6	7.7
ISS 16-24	7.7	7.7
ISS 25-40	7	7.1
ISS 41-75	4.5	5
Hospital length of stay		
Total bed days	602	798.6
Mean overall	7.9	9.3
ISS <13	12.6	17.3
ISS 13-15	3.6	7
ISS 16-24	11	9.2
ISS 25-40	3.5	5.8
ISS 41-75	1	1.2
ICU length of stay		
ICU total bed days (number of ICU admissions)	92 (34)	85.1 (29.2)
Mean overall	2.7	2.9
ISS <13	2.8	2.4
ISS 13-15	3.5	5.6
ISS 16-24	3.5	2.5
ISS 25-40	1.4	2.9
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	46 (16)	20.6 (8.8)
Mean overall	2.9	2.3
ISS <13	3	2.6
ISS 13-15	2	3.7
ISS 16-24	10	2.6
ISS 25-40	2.8	1.9
ISS 41-75	1	1.3

Table 40: Trauma data profile, Gosford Hospital

Description	Facility	Peer
Total admissions	54	86.3
Mean monthly admissions	4.5	7.2
Case fatality rate (ISS >12 excl. traumatic DOA)	8.7%	6.3%
Sex		
Male / Female	38 / 16	62 / 25
Age ranges		
Mean age	58.3	54.4
0-4	0 (0%)	1.1 (1.3%)
5-9	1 (1.9%)	0.9 (1%)
10-14	2 (3.7%)	1.1 (1.3%)
15-19	3 (5.6%)	4 (4.6%)
20-24	4 (7.4%)	6 (7%)
25-29	3 (5.6%)	4.7 (5.4%)
30-34	0 (0%)	3.5 (4.1%)
35-39	0 (0%)	2.9 (3.4%)
40-44	2 (3.7%)	5.2 (6%)
45-49	5 (9.3%)	6.2 (7.2%)
50-54	1 (1.9%)	6.7 (7.8%)
55-59	4 (7.4%)	5.7 (6.6%)
60-64	3 (5.6%)	5.6 (6.5%)
65-69	3 (5.6%)	4.8 (5.6%)
70-74	5 (9.3%)	5.8 (6.7%)
75-79	2 (3.7%)	5 (5.8%)
80-84	4 (7.4%)	7.6 (8.8%)
85+	12 (22.2%)	9.5 (11%)
Injury Severity Score		
Mean ISS	15.2	17.3
ISS <13	8 (14.8%)	14.5 (16.9%)
ISS 13-15	23 (42.6%)	22.6 (26.3%)
ISS 16-24	16 (29.6%)	32.7 (38%)
ISS 25-40	7 (13%)	15.1 (17.6%)
ISS 41-75	0 (0%)	1.1 (1.3%)
Mechanism of injury		
Assault	1 (1.9%)	5.5 (6.4%)
Falls	30 (55.6%)	34.2 (39.6%)
Falls ≥65 years (% of all mechanisms of injury for ≥65 years)	21 (80.8%)	23.1 (70.6%)
Road trauma	10 (18.5%)	25.1 (29.1%)
Other transport incident	4 (7.4%)	12.1 (14%)
All other injuries	9 (16.7%)	9.4 (10.9%)

Description	Facility	Peer
Injury type		
Blunt	52 (96.3%)	81 (93.9%)
Penetrating	2 (3.7%)	4.1 (4.8%)
Unknown	0 (0%)	1.2 (1.4%)
Admission type		
Direct admission	54 (100%)	70.3 (81.5%)
Transfer in	0 (0%)	15.2 (17.6%)
Unknown	0 (0%)	0.8 (0.9%)
Arrival modes		
Ambulance	49 (90.7%)	68.5 (79.4%)
Helicopter	0 (0%)	4.5 (5.2%)
Other (private vehicle, fixed wing aircraft, unknown)	5 (9.3%)	13.3 (15.4%)
Revised Trauma Score		
Mean overall	7.6	7.5
ISS <13	7.7	7.6
ISS 13-15	7.8	7.7
ISS 16-24	7.5	7.7
ISS 25-40	6.8	7.1
Hospital length of stay		
Total bed days	371	798.6
Mean overall	6.9	9.3
ISS <13	8.1	17.3
ISS 13-15	7.9	7
ISS 16-24	6.2	9.2
ISS 25-40	3.6	5.8
ICU length of stay		
ICU total bed days (number of ICU admissions)	16 (3)	85.1 (29.2)
Mean overall	5.3	2.9
ISS <13	7	2.4
ISS 16-24	2	2.5
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	0 (0)	20.6 (8.8)
Mean overall	0	2.3
ISS <13	0	2.6
ISS 13-15	0	3.7
ISS 16-24	0	2.6
ISS 25-40	0	1.9
ISS 41-75	0	1.3

Table 41: Trauma data profile, Lismore Base Hospital

Description	Facility	Peer
Total admissions	78	86.3
Mean monthly admissions	6.5	7.2
Case fatality rate (ISS >12 excl. traumatic DOA)	13.7%	6.3%
Sex		
Male / Female	55 / 23	62 / 25
Age ranges		
Mean age	56.9	54.4
0-4	1 (1.3%)	1.1 (1.3%)
5-9	1 (1.3%)	0.9 (1%)
10-14	1 (1.3%)	1.1 (1.3%)
15-19	3 (3.8%)	4 (4.6%)
20-24	4 (5.1%)	6 (7%)
25-29	3 (3.8%)	4.7 (5.4%)
30-34	1 (1.3%)	3.5 (4.1%)
35-39	2 (2.6%)	2.9 (3.4%)
40-44	10 (12.8%)	5.2 (6%)
45-49	3 (3.8%)	6.2 (7.2%)
50-54	4 (5.1%)	6.7 (7.8%)
55-59	6 (7.7%)	5.7 (6.6%)
60-64	4 (5.1%)	5.6 (6.5%)
65-69	9 (11.5%)	4.8 (5.6%)
70-74	4 (5.1%)	5.8 (6.7%)
75-79	7 (9%)	5 (5.8%)
80-84	5 (6.4%)	7.6 (8.8%)
85+	10 (12.8%)	9.5 (11%)
Injury Severity Score		
Mean ISS	19.9	17.3
ISS <13	4 (5.2%)	14.5 (16.9%)
ISS 13-15	18 (23.4%)	22.6 (26.3%)
ISS 16-24	35 (45.5%)	32.7 (38%)
ISS 25-40	17 (22.1%)	15.1 (17.6%)
ISS 41-75	3 (3.9%)	1.1 (1.3%)
Mechanism of injury		
Assault	2 (2.6%)	5.5 (6.4%)
Falls	28 (35.9%)	34.2 (39.6%)
Falls ≥65 years (% of all mechanisms of injury for ≥65 years)	20 (57.1%)	23.1 (70.6%)
Road trauma	28 (35.9%)	25.1 (29.1%)
Other transport incident	8 (10.3%)	12.1 (14%)
All other injuries	12 (15.4%)	9.4 (10.9%)

Description	Facility	Peer
Injury type		
Blunt	72 (92.3%)	81 (93.9%)
Penetrating	4 (5.1%)	4.1 (4.8%)
Unknown	2 (2.6%)	1.2 (1.4%)
Admission type		
Direct admission	70 (89.7%)	70.3 (81.5%)
Transfer in	8 (10.3%)	15.2 (17.6%)
Unknown	0 (0%)	0.8 (0.9%)
Arrival modes		
Ambulance	61 (78.2%)	68.5 (79.4%)
Helicopter	10 (12.8%)	4.5 (5.2%)
Other (private vehicle, fixed wing aircraft, unknown)	7 (9%)	13.3 (15.4%)
Revised Trauma Score		
Mean overall	7.2	7.5
ISS <13	5.9	7.6
ISS 13-15	7.6	7.7
ISS 16-24	7.7	7.7
ISS 25-40	6.5	7.1
ISS 41-75	5.2	5
Hospital length of stay		
Total bed days	451	798.6
Mean overall	5.8	9.3
ISS <13	6.2	17.3
ISS 13-15	10.9	7
ISS 16-24	4.2	9.2
ISS 25-40	4.6	5.8
ISS 41-75	1	1.2
ICU length of stay		
ICU total bed days (number of ICU admissions)	125 (32)	85.1 (29.2)
Mean overall	3.9	2.9
ISS <13	3.7	2.4
ISS 13-15	6	5.6
ISS 16-24	1.8	2.5
ISS 25-40	3.2	2.9
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	19 (12)	20.6 (8.8)
Mean overall	1.6	2.3
ISS <13	1	2.6
ISS 13-15	2	3.7
ISS 16-24	3	2.6
ISS 25-40	1.6	1.9
ISS 41-75	1	1.3

Table 42: Trauma data profile, Nepean Hospital

Description	Facility	Peer
Total admissions	120	86.3
Mean monthly admissions	10	7.2
Case fatality rate (ISS >12 excl. traumatic DOA)	9%	6.3%
Sex		
Male / Female	72 / 48	62 / 25
Age ranges		
Mean age	60.2	54.4
0-4	1 (0.8%)	1.1 (1.3%)
5-9	0 (0%)	0.9 (1%)
10-14	0 (0%)	1.1 (1.3%)
15-19	3 (2.5%)	4 (4.6%)
20-24	7 (5.8%)	6 (7%)
25-29	6 (5%)	4.7 (5.4%)
30-34	6 (5%)	3.5 (4.1%)
35-39	1 (0.8%)	2.9 (3.4%)
40-44	9 (7.5%)	5.2 (6%)
45-49	5 (4.2%)	6.2 (7.2%)
50-54	10 (8.3%)	6.7 (7.8%)
55-59	3 (2.5%)	5.7 (6.6%)
60-64	4 (3.3%)	5.6 (6.5%)
65-69	8 (6.7%)	4.8 (5.6%)
70-74	16 (13.3%)	5.8 (6.7%)
75-79	11 (9.2%)	5 (5.8%)
80-84	22 (18.3%)	7.6 (8.8%)
85+	8 (6.7%)	9.5 (11%)
Injury Severity Score		
Mean ISS	13.3	17.3
ISS <13	53 (44.2%)	14.5 (16.9%)
ISS 13-15	27 (22.5%)	22.6 (26.3%)
ISS 16-24	27 (22.5%)	32.7 (38%)
ISS 25-40	13 (10.8%)	15.1 (17.6%)
ISS 41-75	0 (0%)	1.1 (1.3%)
Mechanism of injury		
Assault	12 (10%)	5.5 (6.4%)
Falls	72 (60%)	34.2 (39.6%)
Falls ≥65 years (% of all mechanisms of injury for ≥65 years)	59 (90.8%)	23.1 (70.6%)
Road trauma	14 (11.7%)	25.1 (29.1%)
Other transport incident	11 (9.2%)	12.1 (14%)
All other injuries	11 (9.2%)	9.4 (10.9%)

Description	Facility	Peer
Injury type		
Blunt	113 (94.2%)	81 (93.9%)
Penetrating	7 (5.8%)	4.1 (4.8%)
Unknown	0 (0%)	1.2 (1.4%)
Admission type		
Direct admission	93 (77.5%)	70.3 (81.5%)
Transfer in	26 (21.7%)	15.2 (17.6%)
Unknown	1 (0.8%)	0.8 (0.9%)
Arrival modes		
Ambulance	98 (81.7%)	68.5 (79.4%)
Helicopter	0 (0%)	4.5 (5.2%)
Other (private vehicle, fixed wing aircraft, unknown)	22 (18.3%)	13.3 (15.4%)
Revised Trauma Score		
Mean overall	7.7	7.5
ISS <13	7.8	7.6
ISS 13-15	7.8	7.7
ISS 16-24	7.8	7.7
ISS 25-40	7.3	7.1
Hospital length of stay		
Total bed days	2730	798.6
Mean overall	22.8	9.3
ISS <13	34	17.3
ISS 13-15	15.7	7
ISS 16-24	8.3	9.2
ISS 25-40	21.6	5.8
ICU length of stay		
ICU total bed days (number of ICU admissions)	253 (73)	85.1 (29.2)
Mean overall	3.5	2.9
ISS <13	2.5	2.4
ISS 13-15	6.6	5.6
ISS 16-24	2	2.5
ISS 25-40	7.1	2.9
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	0 (0)	20.6 (8.8)
Mean overall	0	2.3
ISS <13	0	2.6
ISS 13-15	0	3.7
ISS 16-24	0	2.6
ISS 25-40	0	1.9
ISS 41-75	0	1.3

Table 43: Trauma data profile, Orange Base Hospital

Description	Facility	Peer
Total admissions	111	86.3
Mean monthly admissions	9.2	7.2
Case fatality rate (ISS >12 excl. traumatic DOA)	3.3%	6.3%
Sex		
Male / Female	93 / 18	62 / 25
Age ranges		
Mean age	49.9	54.4
0-4	2 (1.8%)	1.1 (1.3%)
5-9	1 (0.9%)	0.9 (1%)
10-14	4 (3.6%)	1.1 (1.3%)
15-19	5 (4.5%)	4 (4.6%)
20-24	10 (9%)	6 (7%)
25-29	3 (2.7%)	4.7 (5.4%)
30-34	4 (3.6%)	3.5 (4.1%)
35-39	5 (4.5%)	2.9 (3.4%)
40-44	8 (7.2%)	5.2 (6%)
45-49	11 (9.9%)	6.2 (7.2%)
50-54	13 (11.7%)	6.7 (7.8%)
55-59	10 (9%)	5.7 (6.6%)
60-64	8 (7.2%)	5.6 (6.5%)
65-69	2 (1.8%)	4.8 (5.6%)
70-74	5 (4.5%)	5.8 (6.7%)
75-79	4 (3.6%)	5 (5.8%)
80-84	7 (6.3%)	7.6 (8.8%)
85+	9 (8.1%)	9.5 (11%)
Injury Severity Score		
Mean ISS	16.8	17.3
ISS <13	19 (17.4%)	14.5 (16.9%)
ISS 13-15	32 (29.4%)	22.6 (26.3%)
ISS 16-24	48 (44%)	32.7 (38%)
ISS 25-40	8 (7.3%)	15.1 (17.6%)
ISS 41-75	2 (1.8%)	1.1 (1.3%)
Mechanism of injury		
Assault	4 (3.6%)	5.5 (6.4%)
Falls	30 (27%)	34.2 (39.6%)
Falls ≥65 years (% of all mechanisms of injury for ≥65 years)	17 (63%)	23.1 (70.6%)
Road trauma	39 (35.1%)	25.1 (29.1%)
Other transport incident	25 (22.5%)	12.1 (14%)
All other injuries	13 (11.7%)	9.4 (10.9%)

Description	Facility	Peer
Injury type		
Blunt	108 (97.3%)	81 (93.9%)
Penetrating	3 (2.7%)	4.1 (4.8%)
Unknown	0 (0%)	1.2 (1.4%)
Admission type		
Direct admission	77 (69.4%)	70.3 (81.5%)
Transfer in	28 (25.2%)	15.2 (17.6%)
Unknown	6 (5.4%)	0.8 (0.9%)
Arrival modes		
Ambulance	67 (60.4%)	68.5 (79.4%)
Helicopter	18 (16.2%)	4.5 (5.2%)
Other (private vehicle, fixed wing aircraft, unknown)	26 (23.4%)	13.3 (15.4%)
Revised Trauma Score		
Mean overall	7.8	7.5
ISS <13	7.6	7.6
ISS 13-15	7.8	7.7
ISS 16-24	7.8	7.7
ISS 25-40	7.7	7.1
ISS 41-75	7.8	5
Hospital length of stay		
Total bed days	709	798.6
Mean overall	6.5	9.3
ISS <13	6.4	17.3
ISS 13-15	5.5	7
ISS 16-24	8.1	9.2
ISS 25-40	2.9	5.8
ISS 41-75	1	1.2
ICU length of stay		
ICU total bed days (number of ICU admissions)	65 (31)	85.1 (29.2)
Mean overall	2.1	2.9
ISS <13	1.9	2.4
ISS 13-15	1	5.6
ISS 16-24	2.8	2.5
ISS 25-40	1	2.9
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	10 (3)	20.6 (8.8)
Mean overall	3.3	2.3
ISS <13	3	2.6
ISS 13-15	0	3.7
ISS 16-24	3	2.6
ISS 25-40	0	1.9
ISS 41-75	0	1.3

Table 44: Trauma data profile, Port Macquarie Base Hospital

Description	Facility	Peer
Total admissions	67	86.3
Mean monthly admissions	5.6	7.2
Case fatality rate (ISS >12 excl. traumatic DOA)	1.8%	6.3%
Sex		
Male / Female	47 / 20	62 / 25
Age ranges		
Mean age	47.3	54.4
0-4	1 (1.5%)	1.1 (1.3%)
5-9	1 (1.5%)	0.9 (1%)
10-14	1 (1.5%)	1.1 (1.3%)
15-19	4 (6%)	4 (4.6%)
20-24	8 (11.9%)	6 (7%)
25-29	3 (4.5%)	4.7 (5.4%)
30-34	6 (9%)	3.5 (4.1%)
35-39	3 (4.5%)	2.9 (3.4%)
40-44	4 (6%)	5.2 (6%)
45-49	7 (10.4%)	6.2 (7.2%)
50-54	2 (3%)	6.7 (7.8%)
55-59	1 (1.5%)	5.7 (6.6%)
60-64	5 (7.5%)	5.6 (6.5%)
65-69	5 (7.5%)	4.8 (5.6%)
70-74	9 (13.4%)	5.8 (6.7%)
75-79	2 (3%)	5 (5.8%)
80-84	4 (6%)	7.6 (8.8%)
85+	1 (1.5%)	9.5 (11%)
Injury Severity Score		
Mean ISS	19	17.3
ISS <13	10 (14.9%)	14.5 (16.9%)
ISS 13-15	16 (23.9%)	22.6 (26.3%)
ISS 16-24	25 (37.3%)	32.7 (38%)
ISS 25-40	15 (22.4%)	15.1 (17.6%)
ISS 41-75	1 (1.5%)	1.1 (1.3%)
Mechanism of injury		
Assault	5 (7.5%)	5.5 (6.4%)
Falls	19 (28.4%)	34.2 (39.6%)
Falls ≥65 years (% of all mechanisms of injury for ≥65 years)	8 (38.1%)	23.1 (70.6%)
Road trauma	27 (40.3%)	25.1 (29.1%)
Other transport incident	9 (13.4%)	12.1 (14%)
All other injuries	7 (10.4%)	9.4 (10.9%)

Description	Facility	Peer
Injury type		
Blunt	56 (83.6%)	81 (93.9%)
Penetrating	8 (11.9%)	4.1 (4.8%)
Unknown	3 (4.5%)	1.2 (1.4%)
Admission type		
Direct admission	62 (92.5%)	70.3 (81.5%)
Transfer in	5 (7.5%)	15.2 (17.6%)
Unknown	0 (0%)	0.8 (0.9%)
Arrival modes		
Ambulance	60 (89.6%)	68.5 (79.4%)
Helicopter	1 (1.5%)	4.5 (5.2%)
Other (private vehicle, fixed wing aircraft, unknown)	6 (9%)	13.3 (15.4%)
Revised Trauma Score		
Mean overall	7.6	7.5
ISS <13	7.7	7.6
ISS 13-15	7.8	7.7
ISS 16-24	7.8	7.7
ISS 25-40	7.1	7.1
ISS 41-75	5.7	5
Hospital length of stay		
Total bed days	265	798.6
Mean overall	4	9.3
ISS <13	4.2	17.3
ISS 13-15	5.9	7
ISS 16-24	3.8	9.2
ISS 25-40	2.1	5.8
ISS 41-75	1	1.2
ICU length of stay		
ICU total bed days (number of ICU admissions)	57 (30)	85.1 (29.2)
Mean overall	1.9	2.9
ISS <13	1.4	2.4
ISS 13-15	3	5.6
ISS 16-24	2.3	2.5
ISS 25-40	1.6	2.9
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	16 (11)	20.6 (8.8)
Mean overall	1.5	2.3
ISS <13	2	2.6
ISS 13-15	1.5	3.7
ISS 16-24	2	2.6
ISS 25-40	1	1.9
ISS 41-75	1	1.3

Table 45: Trauma data profile, Tamworth Base Hospital

Description	Facility	Peer
Total admissions	91	86.3
Mean monthly admissions	7.6	7.2
Case fatality rate (ISS >12 excl. traumatic DOA)	2.8%	6.3%
Sex		
Male / Female	68 / 23	62 / 25
Age ranges		
Mean age	49.9	54.4
0-4	2 (2.2%)	1.1 (1.3%)
5-9	1 (1.1%)	0.9 (1%)
10-14	1 (1.1%)	1.1 (1.3%)
15-19	5 (5.5%)	4 (4.6%)
20-24	7 (7.7%)	6 (7%)
25-29	5 (5.5%)	4.7 (5.4%)
30-34	7 (7.7%)	3.5 (4.1%)
35-39	5 (5.5%)	2.9 (3.4%)
40-44	3 (3.3%)	5.2 (6%)
45-49	2 (2.2%)	6.2 (7.2%)
50-54	11 (12.1%)	6.7 (7.8%)
55-59	8 (8.8%)	5.7 (6.6%)
60-64	9 (9.9%)	5.6 (6.5%)
65-69	5 (5.5%)	4.8 (5.6%)
70-74	5 (5.5%)	5.8 (6.7%)
75-79	5 (5.5%)	5 (5.8%)
80-84	4 (4.4%)	7.6 (8.8%)
85+	6 (6.6%)	9.5 (11%)
Injury Severity Score		
Mean ISS	16.8	17.3
ISS <13	19 (20.9%)	14.5 (16.9%)
ISS 13-15	23 (25.3%)	22.6 (26.3%)
ISS 16-24	32 (35.2%)	32.7 (38%)
ISS 25-40	17 (18.7%)	15.1 (17.6%)
ISS 41-75	0 (0%)	1.1 (1.3%)
Mechanism of injury		
Assault	6 (6.6%)	5.5 (6.4%)
Falls	23 (25.3%)	34.2 (39.6%)
Falls ≥65 years (% of all mechanisms of injury for ≥65 years)	14 (56%)	23.1 (70.6%)
Road trauma	24 (26.4%)	25.1 (29.1%)
Other transport incident	22 (24.2%)	12.1 (14%)
All other injuries	16 (17.6%)	9.4 (10.9%)

Description	Facility	Peer
Injury type		
Blunt	80 (87.9%)	81 (93.9%)
Penetrating	7 (7.7%)	4.1 (4.8%)
Unknown	4 (4.4%)	1.2 (1.4%)
Admission type		
Direct admission	70 (76.9%)	70.3 (81.5%)
Transfer in	20 (22%)	15.2 (17.6%)
Unknown	1 (1.1%)	0.8 (0.9%)
Arrival modes		
Ambulance	61 (67%)	68.5 (79.4%)
Helicopter	12 (13.2%)	4.5 (5.2%)
Other (private vehicle, fixed wing aircraft, unknown)	18 (19.8%)	13.3 (15.4%)
Revised Trauma Score		
Mean overall	7.4	7.5
ISS <13	7.5	7.6
ISS 13-15	7.5	7.7
ISS 16-24	7.5	7.7
ISS 25-40	7.2	7.1
Hospital length of stay		
Total bed days	1037	798.6
Mean overall	11.4	9.3
ISS <13	7.9	17.3
ISS 13-15	5.6	7
ISS 16-24	20.7	9.2
ISS 25-40	5.8	5.8
ICU length of stay		
ICU total bed days (number of ICU admissions)	120 (41)	85.1 (29.2)
Mean overall	2.9	2.9
ISS <13	1.6	2.4
ISS 13-15	14.7	5.6
ISS 16-24	2	2.5
ISS 25-40	2.6	2.9
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	42 (12)	20.6 (8.8)
Mean overall	3.5	2.3
ISS <13	2.3	2.6
ISS 13-15	16	3.7
ISS 16-24	2	2.6
ISS 25-40	2.7	1.9
ISS 41-75	0	1.3

Table 46: Trauma data profile, The Tweed Hospital

Description	Facility	Peer
Total admissions	46	86.3
Mean monthly admissions	3.8	7.2
Case fatality rate (ISS >12 excl. traumatic DOA)	5.3%	6.3%
Sex		
Male / Female	27 / 19	62 / 25
Age ranges		
Mean age	55.7	54.4
0-4	1 (2.2%)	1.1 (1.3%)
5-9	1 (2.2%)	0.9 (1%)
10-14	0 (0%)	1.1 (1.3%)
15-19	4 (8.7%)	4 (4.6%)
20-24	4 (8.7%)	6 (7%)
25-29	0 (0%)	4.7 (5.4%)
30-34	1 (2.2%)	3.5 (4.1%)
35-39	1 (2.2%)	2.9 (3.4%)
40-44	2 (4.3%)	5.2 (6%)
45-49	4 (8.7%)	6.2 (7.2%)
50-54	2 (4.3%)	6.7 (7.8%)
55-59	5 (10.9%)	5.7 (6.6%)
60-64	1 (2.2%)	5.6 (6.5%)
65-69	3 (6.5%)	4.8 (5.6%)
70-74	2 (4.3%)	5.8 (6.7%)
75-79	4 (8.7%)	5 (5.8%)
80-84	4 (8.7%)	7.6 (8.8%)
85+	7 (15.2%)	9.5 (11%)
Injury Severity Score		
Mean ISS	18.7	17.3
ISS <13	7 (15.2%)	14.5 (16.9%)
ISS 13-15	6 (13%)	22.6 (26.3%)
ISS 16-24	21 (45.7%)	32.7 (38%)
ISS 25-40	11 (23.9%)	15.1 (17.6%)
ISS 41-75	1 (2.2%)	1.1 (1.3%)
Mechanism of injury		
Assault	1 (2.2%)	5.5 (6.4%)
Falls	18 (39.1%)	34.2 (39.6%)
Falls ≥65 years (% of all mechanisms of injury for ≥65 years)	13 (65%)	23.1 (70.6%)
Road trauma	18 (39.1%)	25.1 (29.1%)
Other transport incident	4 (8.7%)	12.1 (14%)
All other injuries	5 (10.9%)	9.4 (10.9%)

Description	Facility	Peer
Injury type		
Blunt	44 (95.7%)	81 (93.9%)
Penetrating	0 (0%)	4.1 (4.8%)
Unknown	2 (4.3%)	1.2 (1.4%)
Admission type		
Direct admission	41 (89.1%)	70.3 (81.5%)
Transfer in	5 (10.9%)	15.2 (17.6%)
Unknown	0 (0%)	0.8 (0.9%)
Arrival modes		
Ambulance	41 (89.1%)	68.5 (79.4%)
Helicopter	0 (0%)	4.5 (5.2%)
Other (private vehicle, fixed wing aircraft, unknown)	5 (10.9%)	13.3 (15.4%)
Revised Trauma Score		
Mean overall	7.5	7.5
ISS <13	7.8	7.6
ISS 13-15	7.8	7.7
ISS 16-24	7.4	7.7
ISS 25-40	7.5	7.1
ISS 41-75	3.4	5
Hospital length of stay		
Total bed days	552	798.6
Mean overall	12	9.3
ISS <13	7.9	17.3
ISS 13-15	5	7
ISS 16-24	20.6	9.2
ISS 25-40	2.8	5.8
ISS 41-75	3	1.2
ICU length of stay		
ICU total bed days (number of ICU admissions)	35 (14)	85.1 (29.2)
Mean overall	2.5	2.9
ISS <13	2.3	2.4
ISS 13-15	2	5.6
ISS 16-24	4.5	2.5
ISS 25-40	2	2.9
ISS 41-75	2	2
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	13 (7)	20.6 (8.8)
Mean overall	1.9	2.3
ISS <13	2	2.6
ISS 13-15	0	3.7
ISS 16-24	1	2.6
ISS 25-40	2	1.9
ISS 41-75	3	1.3

Table 47: Trauma data profile, Wagga Wagga Base Hospital

Description	Facility	Peer
Total admissions	81	86.3
Mean monthly admissions	6.8	7.2
Case fatality rate (ISS >12 excl. traumatic DOA)	2.7%	6.3%
Sex		
Male / Female	54 / 27	62 / 25
Age ranges		
Mean age	51.1	54.4
0-4	2 (2.5%)	1.1 (1.3%)
5-9	2 (2.5%)	0.9 (1%)
10-14	1 (1.2%)	1.1 (1.3%)
15-19	3 (3.7%)	4 (4.6%)
20-24	4 (4.9%)	6 (7%)
25-29	9 (11.1%)	4.7 (5.4%)
30-34	4 (4.9%)	3.5 (4.1%)
35-39	1 (1.2%)	2.9 (3.4%)
40-44	6 (7.4%)	5.2 (6%)
45-49	5 (6.2%)	6.2 (7.2%)
50-54	7 (8.6%)	6.7 (7.8%)
55-59	9 (11.1%)	5.7 (6.6%)
60-64	6 (7.4%)	5.6 (6.5%)
65-69	2 (2.5%)	4.8 (5.6%)
70-74	3 (3.7%)	5.8 (6.7%)
75-79	3 (3.7%)	5 (5.8%)
80-84	2 (2.5%)	7.6 (8.8%)
85+	12 (14.8%)	9.5 (11%)
Injury Severity Score		
Mean ISS	17	17.3
ISS <13	7 (8.6%)	14.5 (16.9%)
ISS 13-15	30 (37%)	22.6 (26.3%)
ISS 16-24	31 (38.3%)	32.7 (38%)
ISS 25-40	13 (16%)	15.1 (17.6%)
ISS 41-75	0 (0%)	1.1 (1.3%)
Mechanism of injury		
Assault	9 (11.1%)	5.5 (6.4%)
Falls	27 (33.3%)	34.2 (39.6%)
Falls ≥65 years (% of all mechanisms of injury for ≥65 years)	17 (77.3%)	23.1 (70.6%)
Road trauma	24 (29.6%)	25.1 (29.1%)
Other transport incident	16 (19.8%)	12.1 (14%)
All other injuries	5 (6.2%)	9.4 (10.9%)

Description	Facility	Peer
Injury type		
Blunt	80 (98.8%)	81 (93.9%)
Penetrating	1 (1.2%)	4.1 (4.8%)
Unknown	0 (0%)	1.2 (1.4%)
Admission type		
Direct admission	63 (77.8%)	70.3 (81.5%)
Transfer in	18 (22.2%)	15.2 (17.6%)
Unknown	0 (0%)	0.8 (0.9%)
Arrival modes		
Ambulance	63 (77.8%)	68.5 (79.4%)
Helicopter	1 (1.2%)	4.5 (5.2%)
Other (private vehicle, fixed wing aircraft, unknown)	17 (21%)	13.3 (15.4%)
Revised Trauma Score		
Mean overall	7.5	7.5
ISS <13	7.6	7.6
ISS 13-15	7.6	7.7
ISS 16-24	7.8	7.7
ISS 25-40	6.8	7.1
Hospital length of stay		
Total bed days	240	798.6
Mean overall	3	9.3
ISS <13	4.3	17.3
ISS 13-15	2.9	7
ISS 16-24	3.5	9.2
ISS 25-40	1.2	5.8
ICU length of stay		
ICU total bed days (number of ICU admissions)	39 (16)	85.1 (29.2)
Mean overall	2.4	2.9
ISS <13	2.3	2.4
ISS 13-15	2.7	5.6
ISS 16-24	4	2.5
ISS 25-40	1.2	2.9
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	14 (12)	20.6 (8.8)
Mean overall	1.2	2.3
ISS <13	1.5	2.6
ISS 13-15	1.5	3.7
ISS 16-24	1	2.6
ISS 25-40	1	1.9
ISS 41-75	0	1.3

Table 48: Trauma data profile, Wollongong Hospital

Description	Facility	Peer
Total Admissions	139	86.3
Mean monthly admissions	11.6	7.2
Case fatality rate (ISS >12 excl. traumatic DOA)	8.3%	6.3%
Sex		
Male / Female	104 / 35	62 / 25
Age ranges		
Mean age	60.3	54.4
0-4	1 (0.7%)	1.1 (1.3%)
5-9	1 (0.7%)	0.9 (1%)
10-14	0 (0%)	1.1 (1.3%)
15-19	5 (3.6%)	4 (4.6%)
20-24	6 (4.3%)	6 (7%)
25-29	7 (5%)	4.7 (5.4%)
30-34	3 (2.2%)	3.5 (4.1%)
35-39	8 (5.8%)	2.9 (3.4%)
40-44	5 (3.6%)	5.2 (6%)
45-49	14 (10.1%)	6.2 (7.2%)
50-54	8 (5.8%)	6.7 (7.8%)
55-59	7 (5%)	5.7 (6.6%)
60-64	8 (5.8%)	5.6 (6.5%)
65-69	5 (3.6%)	4.8 (5.6%)
70-74	7 (5%)	5.8 (6.7%)
75-79	9 (6.5%)	5 (5.8%)
80-84	22 (15.8%)	7.6 (8.8%)
85+	23 (16.5%)	9.5 (11%)
Injury Severity Score		
Mean ISS	18.3	17.3
ISS <13	5 (3.6%)	14.5 (16.9%)
ISS 13-15	40 (28.8%)	22.6 (26.3%)
ISS 16-24	62 (44.6%)	32.7 (38%)
ISS 25-40	32 (23%)	15.1 (17.6%)
ISS 41-75	0 (0%)	1.1 (1.3%)
Mechanism of injury		
Assault	11 (7.9%)	5.5 (6.4%)
Falls	75 (54%)	34.2 (39.6%)
Falls ≥65 years (% of all mechanisms of injury for ≥65 years)	53 (80.3%)	23.1 (70.6%)
Road trauma	32 (23%)	25.1 (29.1%)
Other transport incident	15 (10.8%)	12.1 (14%)
All other injuries	6 (4.3%)	9.4 (10.9%)

Description	Facility	Peer
Injury type		
Blunt	136 (97.8%)	81 (93.9%)
Penetrating	3 (2.2%)	4.1 (4.8%)
Unknown	0 (0%)	1.2 (1.4%)
Admission type		
Direct admission	100 (71.9%)	70.3 (81.5%)
Transfer in	39 (28.1%)	15.2 (17.6%)
Unknown	0 (0%)	0.8 (0.9%)
Arrival modes		
Ambulance	117 (84.2%)	68.5 (79.4%)
Helicopter	0 (0%)	4.5 (5.2%)
Other (private vehicle, fixed wing aircraft, unknown)	22 (15.8%)	13.3 (15.4%)
Revised Trauma Score		
Mean overall	7.6	7.5
ISS <13	7.8	7.6
ISS 13-15	7.8	7.7
ISS 16-24	7.7	7.7
ISS 25-40	7.2	7.1
Hospital length of stay		
Total bed days	1029	798.6
Mean overall	7.4	9.3
ISS <13	9.4	17.3
ISS 13-15	5.7	7
ISS 16-24	8.6	9.2
ISS 25-40	7	5.8
ICU length of stay		
ICU total bed days (number of ICU admissions)	49 (18)	85.1 (29.2)
Mean overall	2.7	2.9
ISS <13	3.8	2.4
ISS 16-24	2.6	2.5
ISS 25-40	2.1	2.9
Hospital ventilation days		
Total ventilation bed days (number of ventilated cases)	46 (15)	20.6 (8.8)
Mean overall	3.1	2.3
ISS <13	4	2.6
ISS 13-15	0	3.7
ISS 16-24	3.3	2.6
ISS 25-40	2.3	1.9
ISS 41-75	0	1.3

Appendix 5: Calculation of the Injury Severity Score

An ISS is calculated for each patient based on the AIS injury severity classification of their specific injuries. The ISS value ranges from 1-75 and is calculated as:

$$ISS = A^2 + B^2 + C^2$$

Where A, B, and C are the highest AIS severity codes in each of the (up to) three most severely injured ISS body regions.

The six ISS body regions are:

- head or neck
- face
- chest
- abdominal or pelvic contents
- extremities or pelvic girdle
- external.

The following example shows how an ISS is calculated from a set of injuries.

Table 49: ISS calculation example

ISS body region	Injury	AIS severity code	Include in ISS calculation?
Head or neck	Small subdural haematoma	AIS-4	Yes
Chest	Bilateral lung contusion	AIS-4	No
Chest	Bilateral flail chest	AIS-5	Yes
Abdominal or pelvic contents	Superficial spleen laceration	AIS-2	Yes
Extremities or pelvic girdle	Left phalange (little toe) fracture	AIS-1	No

On the basis of the above injuries, the ISS is calculated as:


$$ISS = 4^2 + 5^2 + 2^2$$

$$ISS = 45 \text{ (critical injury)}$$

Appendix 6: Australian Statistical Geography Standard Remoteness Areas

The Australian Statistical Geography Standard (ASGS) Remoteness Areas (RA) is based on the Accessibility and Remoteness Index of Australia which defines locations in terms of remoteness, i.e. the physical distance of a location from the nearest urban centre (access to goods and services) based on population size.

Figure 35: Map of 2016 Remoteness areas for Australia



Source: ABS, <http://www.abs.gov.au/websitedbs/d3310114.nsf/home/remoteness+structure>

For further information on ASGS-RA please see [The Australian Statistical Geography Standard \(ASGS\) Remoteness Structure](#) on the Australian Bureau of Statistics website.