


Osteoporotic refracture prevention


Clinical priorities

Osteoporosis is a chronic disease characterised by reduced bone density and strength. It is associated with a heightened vulnerability to minimal trauma fractures (or fragility fractures). People who have had a minimal trauma fracture are at high risk of refracture.

The *NSW Model of care for osteoporotic refracture prevention* outlines evidence-based care for identifying and managing minimal trauma fractures.



This simplified summary highlights four clinical priority areas to reduce the risk of refracture.



CLINICAL OUTCOMES

- ✓ Improved identification of those requiring refracture prevention services
- ✓ Improved uptake of medication, exercise and calcium intake
- ✓ Fewer preventable fractures
- ✓ Better quality of life

The goal of the Leading Better Value Care osteoporotic refracture prevention service is to identify people who have sustained a minimal trauma fracture and coordinate management of their bone health. This will reduce the risk of a refracture and improve their overall health, wellbeing and quality of life. Refer to *ORP site manual* for an implementation guide against the model of care, including key features and supplementary tools and resources.



Identification and triage

People aged 50+ years with fractures related to osteoporosis should be proactively identified by a designated staff member.

- An active search should be conducted for patients with minimal trauma fractures receiving care in emergency departments, inpatient wards, outpatient clinics and community or primary care settings.
- All imaging involving the spine should be screened by radiology for vertebral compression fractures.



Assessment and diagnosis

Within 16 weeks of a fracture, the patient should receive a thorough, person-centred assessment coordinated by a dedicated clinician. This includes:

- assessment of bone health, consisting of bone mineral density testing (DEXA, QCT) and serum vitamin D
- screening for and assessment of falls risk, using validated tools such as FROP-COM, Berg Balance Scale, Timed Up and Go and repeated sit to stand.
- assessment of future fracture risk using FRAX or Garvan tools
- assessment of medical status, including physical function, comorbidities and mental health
- a diagnosis of osteoporosis or osteopenia by a medical officer where indicated.



Treatment initiation

Access should be coordinated for timely, evidence-based and individualised treatment that is recorded in a personalised management plan. This includes:

- provision of osteoporosis education and self-management support
- initiation of appropriate medical interventions inclusive of bone-sparing medications and supplementary treatments
- prescription of resistance exercise or physical activity
- dietary advice on calcium and protein intake
- coordination of falls prevention interventions
- facilitation of comorbidity management.



Coordination of ongoing care

Patients should be supported to ensure continued treatment, intervention and behaviour change. This should involve community services and their general practitioner. This includes:

- review of progress within six months of intervention with review and modification of management plan
- self-management support to recognise progress and address issues
- a plan for transition to appropriate ongoing community-based care.

Evidence

1. Sanders KM, Watts JJ, Abimanyi-Ochom J, et al. [Osteoporosis Costing NSW & ACT: A Burden of Disease Analysis – 2012 to 2022](#). Sydney: Osteoporosis Australia 2017.

2. Center JR, Bliuc D, Nguyen TV, et al. [Risk of Subsequent Fracture after Low-trauma Fracture in Men and Women](#). JAMA. 2007;297(4):387-94.

Australian Commission on Safety and Quality in Health Care. [Hip Fracture Care Clinical Care Standard](#). September 2016.

Ganda K, Puech M, Chen JS, et al. [Models of Care for the Secondary Prevention of Osteoporotic Fractures: A Systematic Review and Meta-analysis](#). Osteoporos Int. 2013;24(2):393-406.

Nakayama A, Major G, Holliday E, et al. [Evidence of Effectiveness of a Fracture Liaison Service to Reduce the Re-fracture Rate](#). Osteoporos Int. 2016;27 (3):873-9.

NSW Agency for Clinical Innovation. [Model of Care for Osteoporotic Refracture Prevention](#). 2nd edn. Chatswood; ACI:2017.

The Royal Australian College of General Practitioners and Osteoporosis Australia. [Osteoporosis Prevention, Diagnosis and Management in Postmenopausal Women and Men over 50 Years of Age](#). 2nd edn. 2017.